

ORBIFOLD QUASIMAP THEORY

DAEWOONG CHEONG, IONUȚ CIOCAN-FONTANINE, AND BUMSIG KIM

ABSTRACT. We extend to orbifolds the quasimap theory of [8, 12], as well as the genus zero wall-crossing results from [9, 11]. As a consequence, we obtain generalizations of orbifold mirror theorems, in particular, of the mirror theorem for toric orbifolds recently proved independently by Coates, Corti, Iritani, and Tseng [13].

CONTENTS

1. Introduction	2
1.1. Acknowledgments	4
2. The stack of stable quasimaps to an orbifold GIT target	4
2.1. Conventions and notations	4
2.2. Quotients	5
2.3. Stable quasimaps to X	5
2.4. Proof of Theorem 2.3.7	10
2.5. Basic Properties and Variants	13
3. Quasimap Theory	15
3.1. Quasimap Invariants	15
3.2. J^ε -function	17
3.3. Factorization	19
3.4. Unitarity of S^ε	23
3.5. The P -series and Birkhoff factorization of J^ε	23
3.6. ε -wall-crossing	25
3.7. Proof of Theorem 3.6.2	26
4. \mathbb{I} -functions and stacky loop spaces	28
4.1. \mathbb{I} -functions	29
4.2. Stacky loop spaces	30
5. Toric Orbifolds	33
5.1. Set-up	33
5.2. Explicit description of $\mathrm{Hom}_\beta(\mathbb{P}_{a,1}, [W/G])$	34
5.3. The virtual normal bundle	35
References	39

1. INTRODUCTION

Orbifold Gromov-Witten theory was first introduced by Chen and Ruan [7], with an algebraic version due later to Abramovich, Graber, and Vistoli [1].

The theory of ε -stable quasimaps to a large class of GIT quotient targets was developed in [12], generalizing and unifying the earlier works [23, 8, 25, 27]. In the appropriate general context of the theory, the GIT target is a smooth Deligne-Mumford stack (or orbifold), but to keep the technicalities at a reasonable level, [12] worked under the assumption that the GIT quotient is a smooth variety and delegated the orbifold case to subsequent work. In §2 of this paper we formally establish the foundations of quasimap theory for orbifold GIT targets by combining the formalism of [12] with the one developed in [1] for algebraic orbifold Gromov-Witten theory.

Namely, consider a triple (W, G, θ) consisting of an affine variety W over \mathbb{C} , a reductive complex algebraic group G acting on W , and a character θ of G . When there are no strictly semistable points for the linearization induced by θ , we have the *GIT stack quotient* $X := [W^{ss}(\theta)/G] = [W^s(\theta)/G]$. We construct a family, depending on a stability parameter $\varepsilon \in \mathbb{Q}_{>0} \cup \{0+, \infty\}$, of (relative) compactifications of moduli stacks of maps from irreducible twisted marked curves with fixed numerical data to the GIT stack quotient $[W^{ss}/G]$. These compactifications are themselves modular and we prove in Theorem 2.3.7 that they are Deligne-Mumford stacks, proper over the affine quotient. Furthermore, if the singularities of W are at worst lci and the semistable locus W^{ss} is nonsingular, these moduli spaces carry canonical perfect obstruction theories and therefore possess virtual fundamental classes.

Once the moduli spaces with the required properties are constructed, the descendant orbifold ε -quasimap invariants associated to the triple (W, G, θ) are defined as usual via integration against the virtual class of products of tautological psi-classes and of Chen-Ruan cohomology classes pulled-back from the (rigidified) inertia stack of X via the evaluation maps. When $\varepsilon > 2$, they recover the orbifold Gromov-Witten invariants of the DM stack X .

It is natural to seek wall-crossing formulas for the invariants as the stability parameter ε varies. For triples giving GIT quotients which are nonsingular varieties, such wall-crossing formulas in genus zero were conjectured in [9] as equalities of generating series of the invariants after a change of variables. The formulas were proved in [9] in the presence of an action of a torus T on W such that the fixed loci of the induced T -action on $W//G$ have good properties. In section 3 we

provide analogous results for the orbifold case. We describe them informally in this Introduction and refer the reader to Conjecture 3.6.1 and Theorem 3.6.2 for the precise statements. These results may be viewed as generalized mirror theorems for orbifolds.

Recall that the genus zero orbifold Gromov-Witten theory of X is encoded in Givental's Lagrangian cone \mathcal{L}_X and this cone is completely determined by the big J -function $J^\infty(q, t, z)$ of X (the notation reflects that GW theory corresponds to the stability parameter $\varepsilon = \infty$). This is the generating series for all Gromov-Witten invariants with at most one descendent insertion and arbitrary number of primary insertions. It depends on the Novikov variables q , the general Chen-Ruan cohomology class t , and a formal variable z . The t -derivatives of the J -function determine the so-called S -operator S_t^∞ . Conversely, the string equation says that the J -function is obtained from the S -operator: $J^\infty(q, t, z) = S_t^\infty(\mathbb{1}_X)$, where $\mathbb{1}_X$ is the fundamental class. Since the terms of S -operator involve invariants with at least two insertions, it has a direct quasimap analogue S_t^ε for every stability parameter. The wall-crossing formula for S -operators then reads

$$(1.0.1) \quad S_t^\varepsilon(\mathbb{1}_X) = S_{\tau^\varepsilon(t)}^\infty(\mathbb{1}_X),$$

with the ‘‘mirror map’’ $\tau^\varepsilon(t)$ a generating series for primary ε -quasimap invariants with a fundamental class insertion. In particular, the mirror map acquires an enumerative interpretation. Theorem 3.6.2 proves (1.0.1) under the assumption that there is a T -action whose fixed points in X are isolated.

Because some genus zero quasimap moduli spaces require at least two markings, a different construction is needed to extend the J -function to other stability parameters. To this end, recall that the Gromov-Witten big J -function has a well-known expression as a generating series of certain localization residues for the natural \mathbb{C}^* -action on *graph spaces* - moduli spaces of genus zero stable maps with one parametrized component of the domain curve. The graph spaces exist for any stability parameter $\varepsilon \in \mathbb{Q}_{>0} \cup \{0+, \infty\}$ and the corresponding localization residues define the big J^ε -function of the ε -quasimap theory. The wall-crossing/mirror formula for big J -functions is then the statement that $J^\varepsilon(q, t, z)$ is on the Lagrangian cone \mathcal{L}_X for all $\varepsilon \geq 0+$; this is the second part of Theorem 3.6.2. For semi-positive targets (see Definition 3.5.1) this follows from the formula for S -operators. Without the semi-positivity condition a separate proof is needed and our argument requires the additional assumption that the one-dimensional T -orbits in X are also isolated.

If one is primarily interested in determining the orbifold Gromov-Witten invariants of X , a wall-crossing formula becomes useful when the quasimap side can be explicitly computed. While in general the J^ε -functions are equally hard to compute for all stability parameters, in section 4 of the paper we consider a version of moduli spaces of quasimaps, dubbed “stacky loop spaces”, which will typically allow one to find closed formulas for the *small* I -function $I(q, z) := J^{0+}(q, 0, z)$, that is, for the specialization at $t = 0$ of the J -function for the asymptotic stability condition $\varepsilon = 0+$. Following [11], we also introduce a new orbifold “big I -function” $\mathbb{I}_X(q, t, z)$. It is an explicit modification of the small I -function by certain exponential factors and it depends on a parameter t which runs over the part of $H^*(X)$ generated by Chern classes of line bundles associated to characters of G . As in [11], the new \mathbb{I} -function can be viewed as arising from a variant with weighted markings of quasimap theory. The second main result of the paper, Theorem 4.1.2, states that $\mathbb{I}_X(q, t, z)$ is on the Lagrangian cone \mathcal{L}_X whenever the T -action has isolated fixed points and isolated one-dimensional orbits.

As an application of the theory developed in the paper, in section 5 we discuss the case of toric DM stacks. These are the quotients $X = [W^{ss}/G]$ for W a vector space and $G \cong (\mathbb{C}^*)^r$ an algebraic torus. We make Theorem 4.1.2 completely explicit by calculating the small I -function $I(q, z)$ in closed form via stacky loop spaces. This gives a closed form for the “big” \mathbb{I} -function as well. As a result, the Mirror Theorem for toric orbifolds, recently proved by different methods in [13], becomes a special case of Theorem 4.1.2, see Corollary 5.3.4.

1.1. Acknowledgments. The research of D.C. was partially supported by the NRF grant 2007-0093859. The research of I.C.-F. was partially supported by the NSF grant DMS-1305004. The research of B.K. was partially supported by the NRF grant 2014-001824. I.C.-F. thanks KIAS for excellent working conditions during visits when this research was conducted.

2. THE STACK OF STABLE QUASIMAPS TO AN ORBIFOLD GIT TARGET

2.1. Conventions and notations. We work over the field \mathbb{C} . All schemes are locally of finite type over $\text{Spec}\mathbb{C}$ unless otherwise stated. Associated to a DM stack X of finite type over \mathbb{C} , we have the (*cyclo-tomic*) *inertia stack* $I_\mu X$ and its rigidification $\bar{I}_\mu X$ (see [1]). For DM stacks $X, \bar{I}_\mu X, \dots$, we denote by $\underline{X}, \underline{\bar{I}_\mu X}, \dots$ their coarse moduli spaces.

2.2. Quotients. Let W be an irreducible affine variety with a right action of a reductive algebraic group G . Denote by \mathbb{C}_θ the one dimensional G representation space associated to a fixed character θ of G . Denote by W^{ss} (or $W^{ss}(\theta)$) the semistable locus, and by W^s (or $W^s(\theta)$) the stable locus with respect to the linearization $L_\theta := W \times \mathbb{C}_\theta$ (see e.g., [20] for the definitions). There are then four quotients with natural morphisms between them, summarized in the diagram

$$(2.2.1) \quad \begin{array}{ccc} X := [W^{ss}/G] & \hookrightarrow & \mathfrak{X} := [W/G] \\ \downarrow & & \downarrow \\ \underline{X} := W//_\theta G & \longrightarrow & \underline{X}_0 := \text{Spec}\mathbb{C}[W]^G. \end{array}$$

In the top line the brackets denote as usual the stack quotients, and the arrow is the open embedding induced by the inclusion $W^{ss} \subset W$. The GIT quotient $W//_\theta G$ is defined to be

$$\mathbf{Proj} \bigoplus_{m=0}^{\infty} \Gamma(W, L_\theta^m)^G$$

and the bottom arrow is the obvious projective morphism to the affine quotient $\text{Spec}\mathbb{C}[W]^G$. (Note also that the affine quotient coincides with $W//_0 G$.) The stack $[W^{ss}/G]$ will be called the *GIT stack quotient* with respect to θ . The natural vertical morphisms in (2.2.1) are obtained from the fact that principal G -bundles are categorical quotients, since the morphisms

$$W^{ss}(\theta) \rightarrow W//_\theta G, \quad W \rightarrow W//_0 G$$

commute with the G -actions.

In this paper we assume that

$$W^{ss}(\theta) = W^s(\theta)$$

unless otherwise stated so that the GIT stack quotient $X = [W^{ss}/G]$ is a quasi-compact DM-stack. The morphism $X \rightarrow \underline{X}$ is the coarse moduli and is a proper morphism, see e.g. [19]. Therefore X is proper over \underline{X}_0 since \underline{X} is projective over the affine quotient \underline{X}_0 .

2.3. Stable quasimaps to X . Let \mathbf{e} be the least common multiple of the exponents $|\text{Aut}(\bar{p})|$ of automorphism groups $\text{Aut}(\bar{p})$ of all geometric points $\bar{p} \rightarrow X$ of X .

Definition 2.3.1. Let (C, x_1, \dots, x_k) be a k -pointed, genus g twisted curve, see [1, §4] and let $\phi : (C, x_1, \dots, x_k) \rightarrow (\underline{C}, \underline{x}_1, \dots, \underline{x}_k)$ be its coarse moduli space.

A representable morphism $[u]$ from (C, x_1, \dots, x_k) to \mathfrak{X} is called a k -pointed, genus g quasimap to X (alternatively, a θ -quasimap to \mathfrak{X}) if $[u]^{-1}(\mathfrak{X} \setminus X)$ is purely zero-dimensional.

The locus $[u]^{-1}(\mathfrak{X} \setminus X)$ is called the base locus of $[u]$.

The class β of the quasimap is the group homomorphism

$$\beta : \text{Pic}\mathfrak{X} \rightarrow \mathbb{Q}, \quad L \mapsto \deg([u]^*(L)).$$

The rational number $\beta(L_\theta) = \deg([u]^*(L_\theta))$ is called the degree of the quasimap $[u]$.

A quasimap $((C, x_1, \dots, x_k), [u])$ is called prestable if the base locus contains neither marked gerbes nor nodal gerbes of (C, x_1, \dots, x_k) .

Fix a positive rational number ε . A prestable quasimap is called ε -stable if the following two conditions hold:

(1)

$$(2.3.1) \quad \omega_{\underline{C}}(\sum \underline{x}_i) \otimes (\phi_*([u]^* L_\theta^{\otimes e}))^{\varepsilon/e}$$

on the coarse curve \underline{C} is ample.

(2) For every x in C ,

$$\varepsilon l(x) \leq 1,$$

where $l(x)$ is the length at x defined in [12, §7.1].

A prestable quasimap is called $(0+)$ -stable (or simply stable) if it is ε -stable for every small enough positive rational number ε .

A few explanations are in order.

- Throughout this paper, a twisted curve is required to be balanced ([1, §4]).
- The degree of $[u]^*L$ for $L \in \text{Pic}\mathfrak{X}$ is defined using a finite covering of the normalization of C as in [1, §7.2].

Assume that the quasimap $[u]$ is prestable. By the representability of $[u]$ and Lemma 2.1.2 of [1], there is on the coarse moduli space \underline{C} a line bundle whose pullback by ϕ is isomorphic to $[u]^*L^{\otimes e}$ and hence $\deg[u]^*L \in \frac{1}{e}\mathbb{Z}$. In other words, for prestable twisted quasimaps the class β is an element of $\text{Hom}_{\mathbb{Z}}(\text{Pic}\mathfrak{X}, \frac{1}{e}\mathbb{Z})$ and so it has uniformly bounded denominators.

However, note that the definition of the class β as an element of $\text{Hom}_{\mathbb{Z}}(\text{Pic}\mathfrak{X}, \mathbb{Q})$ makes sense for an arbitrary morphism of stacks $[u] : C \rightarrow \mathfrak{X}$ and we will use it later in this generality.

- It is immediate from the definition that a prestable quasimap is ε -stable if and only if

- (1) for every irreducible component C_i of C ,
- (2.3.2) $2g(C_i) - 2 + \# \text{ special points on } C_i + \varepsilon \deg([u]^* L_\theta|_{C_i}) > 0$

and

- (2) $\varepsilon l(x) \leq 1$ for every point x in C .
- For $(g, k) \neq (0, 0)$, a prestable quasimap is a stable twisted map into X if and only if it is an ε -stable quasimap for some $\varepsilon > 1$. For $(g, k) = (0, 0)$, the same is true with $\varepsilon > 2$. For simplicity, a large enough ε will be denoted by ∞ .
 - The base locus of a prestable quasimap $[u]$ will be considered as the reduced scheme.
 - The notions of (prestability, ε -stability) quasimaps over an algebraically closed field of characteristic zero can be identically defined.

From now on we fix $\varepsilon \in [0+, \infty]$. For short, we let $\mathbf{x} = x_1, \dots, x_k$.

Definition 2.3.2. A group homomorphism $\beta \in \text{Hom}_{\mathbb{Z}}(\text{Pic}\mathfrak{X}, \mathbb{Q})$ is called L_θ -effective if it is realized as a finite sum of classes of some quasimaps to X . Such elements form a semigroup with identity 0, denoted by $\text{Eff}(W, G, \theta)$.

Lemma 2.3.3. If $((C, \mathbf{x}), [u])$ is a quasimap of class β , then $\beta(L_\theta) \geq 0$. Moreover, $\beta(L_\theta) = 0$ if and only if $\beta = 0$, if and only if the quasimap is constant (i.e., u is a map into X , factored through an inclusion $B\Gamma \subset X$ of the classifying groupoid $B\Gamma$ of a finite group Γ).

Proof. Consider a finite covering \hat{C} of the normalization of C such that \hat{C} is a (possibly disconnected) nonsingular projective curve. Then the induced map $[\hat{u}] : \hat{C} \rightarrow \mathfrak{X}$ is a union of quasimaps. Now the Lemma follows by applying [12, Lemma 3.2.1] to $[\hat{u}]$. \square

Definition 2.3.4. Let (C, \mathbf{x}) be a family of twisted k -pointed genus g curve over a scheme S , see [1, §4], [26, §1.1]. In particular, the markings $x_i \subset C$ are étale gerbes over S banded by finite cyclic groups. A pair $((C, \mathbf{x}), [u])$ is called a (resp. prestable, ε -stable) quasimap to X over S if $[u]$ is a morphism from C to \mathfrak{X} such that the restrictions to geometric fibers are (resp. prestable, ε -stable) quasimaps in the sense of Definition 2.3.1.

Lemma 2.3.5. Let D be a codimension 1 nonsingular subvariety of a nonsingular variety Y , and let f be a morphism from $Y \setminus D$ to a separated DM stack \mathcal{X} .

Suppose that the coarse moduli level morphism \underline{f} induced by f is extendible to a morphism $Y \rightarrow \underline{\mathcal{X}}$. Then after possibly shrinking Y at

a given closed point p of D (i.e., taking an étale open subset containing p) we have a representable morphism $\tilde{f} : Y_{D,r} \rightarrow \mathcal{X}$ extending f , where $Y_{D,r}$ is the root stack of the effective divisor D with r -twisting for some r (for the definition of root stacks see [6, 1]). The pair $(Y_{D,r}, \tilde{f})$ is unique up to a unique isomorphism, i.e., another one $(Y_{D,r'}, \tilde{f}')$ is isomorphic to the $(Y_{D,r}, \tilde{f})$ up to a unique isomorphism.

Proof. Since this is a local problem in the étale topology, we may assume that $\mathcal{X} = [Z/\Gamma]$ for some affine scheme Z with a finite group Γ action (see [3, Lemma 2.2.3], [19, Proposition 4.2]). Take an analytic neighborhood U of p such that the fundamental group of $U \setminus D$ is isomorphic to \mathbb{Z} . Let $Z_{U \setminus D}$ be the principal Γ -bundle on $U \setminus D$ obtained by pulling back $Z \rightarrow \mathfrak{X}$ via f . We have the monodromy action of \mathbb{Z} on $Z_{U \setminus D}$, which factors through an action of the group μ_r of r -th roots of unity for some positive integer r and induces a monomorphism $\mu_r \rightarrow \Gamma$.

Take an r to 1 covering $U' \rightarrow U$ branched along D , with U' nonsingular. Let $D' \subset U'$ be the reduced divisor corresponding to D . On $U' \setminus D'$, the pull-back bundle $Z_{U' \setminus D'}$ is a trivial Γ -bundle. Choosing a section of it, we get by composition a μ_r -equivariant morphism from $U' \setminus D'$ to Z .

By assumption, the induced map

$$U' \setminus D' \longrightarrow [Z/\Gamma]$$

is extendible to

$$U' \rightarrow U \rightarrow [Z/\Gamma].$$

Since the coarse moduli space $[Z/\Gamma] \rightarrow [Z/\Gamma]$ is proper by [19], after shrinking U' if necessary, there is a branched covering $U'' \rightarrow U'$ along D' and an extension $U'' \rightarrow [Z/\Gamma]$, where U'' is nonsingular. Using a trivialization of the pullback bundle $Z_{U''}$, we obtain a morphism from U'' to Z . This implies that there is a morphism $U' \rightarrow Z$ through which the morphism $U'' \rightarrow Z$ is factored. The morphism $U' \rightarrow Z$ is μ_r -equivariant since it is so generically. It therefore descends to a representable morphism $U_{D,r} := [U'/\mu_r] \rightarrow [Z/\Gamma]$, as claimed in the Lemma. The uniqueness follows from the separatedness of $[Z/\Gamma]$ and the uniqueness of r . \square

We note an immediate consequence of Lemma 2.3.5. If $((C, \mathbf{x}), [u])$ is a prestable quasimap to X , with base locus B (viewed as a reduced subscheme of C), then there is a canonical twisted curve $(C, \mathbf{x} \cup B)$ and a canonical representable morphism $[u]_{\text{reg}} : (C, \mathbf{x} \cup B) \rightarrow X$ which extends $[u]_{|_{C \setminus B}}$. Indeed, the required extension $[u] : \underline{C} \rightarrow \underline{X}$ exists by

the prestable assumption and the fact that $\underline{X} \rightarrow \underline{X}_0$ is projective. As in Lemma 7.1.2 of [12], if β_{reg} denotes the class of $[u]_{\text{reg}}$, then

$$(2.3.3) \quad (\beta - \beta_{\text{reg}})(L_\theta) = \sum_{x \in C} l(x).$$

We will need also the following family version of the above consequence.

Corollary 2.3.6. *Let Δ be a nonsingular curve. Consider a Δ -family of prestable quasimaps $((C, x_1, \dots, x_k), [u])$ to X with base locus B . Then, after possibly shrinking Δ and making an étale base change there is a unique Δ -family of twisted stable maps $(C', x'_1, \dots, x'_k, b'_1, \dots, b'_l, [u]_{\text{reg}})$ into X , together with an isomorphism*

$$\varphi : ((C' \setminus \bigcup \{b'_j\}_j), x'_1, \dots, x'_k) \rightarrow ((C \setminus B), x_1, \dots, x_k)$$

over Δ making $[u] \circ \varphi = [u]_{\text{reg}}$ on $C' \setminus \bigcup \{b'_j\}_j$.

Proof. After shrinking Δ and étale base change, we may assume that the base locus B forms (possibly empty) sections b_j , $j = 1, \dots, l$, of $C \rightarrow \Delta$, disjoint from the markings \mathbf{x} and the nodes in the fibers. Since $[u]$ is a morphism $C \rightarrow \mathfrak{X}$, it induces a morphism $C \rightarrow \underline{X}_0$, which is compatible with $C \setminus B \rightarrow \underline{X}$. Since $\underline{X} \rightarrow \underline{X}_0$ is projective and B is a smooth divisor of C , we may extend $C \setminus B \rightarrow \underline{X}$ at the generic point of every component of B . So, after shrinking Δ again, we may assume that there is an extension $C \rightarrow \underline{X}$. Applying Lemma 2.3.5 concludes the proof. \square

Now we come to the main result of this section. A related statement in the case of one specific GIT target is the main result of [15].

Theorem 2.3.7. *The category fibered in groupoids $Q_{g,k}^\varepsilon(X, \beta)$ of genus g , k -pointed ε -stable quasimaps to X of class β is a DM stack, proper over \underline{X}_0 . Further if W is a locally complete intersection (LCI) variety, then $Q_{g,k}^\varepsilon(X, \beta)$ is equipped with a canonical perfect obstruction theory.*

Precisely speaking, $Q_{g,k}^\varepsilon(X, \beta)$ is a priori a 2-category. An arrow from $(C, \mathbf{x}, [u])$ to $(C', \mathbf{x}', [u'])$ over a morphism $S \rightarrow S'$ between schemes is a pair (φ, α) of a cartesian product $\varphi : C \rightarrow C'$ over $S \rightarrow S'$ preserving the order of markings and a 2-morphism $\alpha : [u] \Rightarrow [u'] \circ \varphi$. A 2-arrow from (φ, α) to (φ', α') is a 2-morphism $\sigma : \varphi \Rightarrow \varphi'$ compatible with α and α' . Since C, C' are DM stacks containing dense open algebraic spaces, the morphisms σ are unique if they exist, by [3, Lemma 4.2.3]. Therefore the 2-category $Q_{g,k}^\varepsilon(X, \beta)$ is equivalent to a category.

In fact, $Q_{g,k}^\varepsilon(X, \beta)$ depends on the pair (\mathfrak{X}, X) , see [12, §4.6]. Therefore, a more precise notation would be $Q_{g,k}^\varepsilon((\mathfrak{X}, X), \beta)$, but we'll only use the latter when needing to emphasize this feature.

When $\varepsilon > 1$ (or $\varepsilon > 2$ if $(g, k) = (0, 0)$), $Q_{g,k}^\varepsilon(X, \beta)$ is nothing but the stack $\mathcal{K}_{g,k}(X, \beta)$ of k -pointed twisted stable maps into X of genus g and class β introduced by Abramovich and Vistoli in [3]. We will also use the notation $Q_{g,k}^\infty(X, \beta) := \mathcal{K}_{g,k}(X, \beta)$ for these moduli stacks.

2.4. Proof of Theorem 2.3.7. In this section we prove Theorem 2.3.7.

2.4.1. Algebraicity. Let $\mathfrak{Qmap}_{g,k}(X, \beta)$ denote the category fibered in groupoids parameterizing k -pointed genus g quasimaps to X of class β (no (pre)stability condition imposed). We will show that $\mathfrak{Qmap}_{g,k}(X, \beta)$ is an Artin stack locally of finite presentation over X_0 .

Let $\mathfrak{M}_{g,k}^{\text{tw}}$ be the category fibered in groupoids of k -pointed genus g twisted curves. It is proven in [26, Theorems 1.9, 1.10] that it is a smooth Artin stack, locally of finite type. Let $\mathfrak{C} \rightarrow \mathfrak{M}_{g,k}^{\text{tw}}$ be the universal curve. We can view $\mathfrak{Qmap}_{g,k}(X, \beta)$ as an open substack of the stack $\text{Hom}_{\mathfrak{M}_{g,k}^{\text{tw}}}(\mathfrak{C}, \mathfrak{X} \times \mathfrak{M}_{g,k}^{\text{tw}})$ whose fiber over a scheme S is the groupoid of 1-morphisms from S -families of k -pointed genus g twisted curves to \mathfrak{X} (for the definition of Hom-stacks see [2, C.1], [22, §2.3]). Hence the desired statement follows from the following

Lemma 2.4.1. *The Hom-stack $\text{Hom}_{\mathfrak{M}_{g,k}^{\text{tw}}}(\mathfrak{C}, \mathfrak{X} \times \mathfrak{M}_{g,k}^{\text{tw}})$ is an Artin stack locally of finite presentation over \mathfrak{C} .*

Proof. Let S be a scheme locally of finite type over \mathfrak{C} , with a smooth surjective morphism $S \rightarrow \mathfrak{M}_{g,k}^{\text{tw}}$. Then the fiber product $S \times_{\mathfrak{M}_{g,k}^{\text{tw}}} \text{Hom}_{\mathfrak{M}_{g,k}^{\text{tw}}}(\mathfrak{C}, \mathfrak{X} \times \mathfrak{M}_{g,k}^{\text{tw}})$ is equivalent to $\text{Hom}_S(\mathfrak{C} \times_{\mathfrak{M}_{g,k}^{\text{tw}}} S, \mathfrak{X} \times S)$, which is an Artin stack locally of finite presentation over S by Proposition 2.11 in [22] since S is an excellent scheme over \mathfrak{C} and $\mathfrak{X} \times S$ is the quotient stack $[W \times S/G]$. Thus, by Lemma C.5 in [2] we conclude the proof. \square

2.4.2. Automorphisms. So far we have shown that $Q_{g,k}^\varepsilon(X, \beta)$ is an Artin stack. By the argument in [12, Proposition 7.1.5], an ε -stable quasimap over a geometric point has no infinitesimal automorphisms. It follows that the diagonal of $Q_{g,k}^\varepsilon(X, \beta)$ is formally unramified, and hence $Q_{g,k}^\varepsilon(X, \beta)$ is Deligne-Mumford.

2.4.3. *Boundedness.* For $d \in \frac{1}{e}\mathbb{Z}_{\geq 0}$, we show that

$$Q_{g,k}^\varepsilon(X, d) := \coprod_{\beta: \beta(L_\theta) = d} Q_{g,k}^\varepsilon(X, \beta),$$

which is locally of finite presentation over \mathbb{C} , is quasi-compact over \mathbb{C} and hence of finite type over \mathbb{C} .

For a fixed twisted curve D and $d \in \frac{1}{e}\mathbb{Z}_{\geq 0}$, $\mathbf{Qmap}_d(D, X)$ will denote the substack of $\mathrm{Hom}(D, \mathfrak{X})$ parameterizing quasimaps $[u]$ from D to X for which $\deg[u]^*(L_\theta) = d$.

(i) *Boundedness of $\mathbf{Qmap}_d(C, X)$:* Let (C, \mathbf{x}) be a twisted curve and choose a projective 1-dimensional variety \tilde{C} with a degree l_C finite flat morphism $\tilde{C} \rightarrow C$ (see [21]). By pullback, there is a natural morphism $\mathrm{Hom}(C, \mathfrak{X}) \rightarrow \mathrm{Hom}(\tilde{C}, \mathfrak{X})$. Since $\tilde{C} \rightarrow C$ is fppf, by the effective descent for principal G -bundles and morphisms we see that

$$\mathrm{Hom}(C, \mathfrak{X}) \cong \mathrm{Hom}(\tilde{C}, \mathfrak{X}) \times_{\mathrm{Hom}(\tilde{C} \times_C \tilde{C}, \mathfrak{X})} \mathrm{Hom}(\tilde{C}, \mathfrak{X}).$$

On the other hand, by Theorem 3.2.4 of [12] (applied to the normalization of \tilde{C}) the moduli stack $\mathbf{Qmap}_{l_C d}(\tilde{C}, X)$ is quasi-compact. It follows that the stack $\mathbf{Qmap}_d(C, X)$ is quasi-compact.

(ii) *Boundedness of the domain curves:* The boundedness of topological types of coarse moduli spaces of possible domain curves follows from (2.3.2) and hence we obtain the boundedness of the possible domain twisted curves.

Now the quasi-compactness of $Q_{g,k}^\varepsilon(X, d)$ follows from (i) and (ii): First, for each $(C, \mathbf{x}) \in \mathfrak{M}_{g,k}^{\mathrm{tw}}(\mathbb{C})$, using (i), we take a finite type scheme $U_{(C, \mathbf{x})}$ smooth over $Q_{g,k}^\varepsilon(X, d)$ containing all quasimaps with the domain curve (C, \mathbf{x}) . Next, by (ii) there is a finite collection $\{U_{(C, \mathbf{x})}\}_{(C, \mathbf{x})}$ surjectively covering $Q_{g,k}^\varepsilon(X, d)$.

Remark 2.4.2. The boundedness of $Q_{g,k}^\varepsilon(X, d)$ implies that there are only finite many $\beta \in \mathrm{Eff}(W, G, \theta)$ with $\beta(L_\theta) = d$.

2.4.4. *Properness.* To prove the properness of $Q_{g,k}^\varepsilon(X, \beta) \rightarrow \underline{X}_0$, it is enough to check the valuative criterion with discrete valuation rings since the stack is already shown to be of finite type over \mathbb{C} . It is straightforward to see that the valuative criterion for separatedness follows from an argument identical to the one given in [8, §4.1].

Once we know the existence of $[u]_{\mathrm{reg}}$ from Corollary 2.3.6, the valuative criterion of properness can be checked by the proof of Theorem 7.1.6 in [12]. The argument requires the properness of the moduli stack $\mathcal{K}_{g,n}(X, \beta)$ of twisted stable maps, which is due to Abramovich and Vistoli [3]. There is a slight modification as follows. In the proof of [12,

Theorem 7.1.6], we need to contract unstable rational tails attached to the central fiber of a completed twisted curve \widehat{C} over a nonsingular curve $(\Delta, 0)$. Those rational tails might have stacky nodal points. When we contract such unstable rational tails, we remove the stack structure of such nodes to make the rational tails (-1) -curves and then we contract those (-1) -curves.

2.4.5. *Obstruction Theory.* This part is identical to the corresponding one in [12]. If we write

$$\mathfrak{Bun}_G^{\text{tw}} := \text{Hom}_{\mathfrak{M}_{g,k}^{\text{tw}}}(\mathfrak{C}, BG \times \mathfrak{M}_{g,k}^{\text{tw}})$$

which is smooth over $\mathfrak{M}_{g,k}^{\text{tw}}$ since the obstruction vanishes (see, e.g., [12, Proposition 2.1.1] for the vanishing of the obstruction), there is a natural forgetful morphism $\sigma : Q_{g,k}^\varepsilon(X, \beta) \rightarrow \mathfrak{Bun}_G^{\text{tw}}$. We describe a canonical σ -relative obstruction theory.

Let $\pi : \mathcal{C} \rightarrow Q_{g,k}^\varepsilon(X, \beta)$ be the universal curve. The universal morphism $[u] : \mathcal{C} \rightarrow \mathfrak{X} = [W/G]$ induces the universal principal G -bundle $\mathcal{P} \rightarrow \mathcal{C}$ by the pullback of the principal G -bundle $W \rightarrow [W/G]$ (here \mathcal{P} is an algebraic space over \mathbb{C} since $[u]$ is representable). In turn, this determines the universal fiber bundle

$$\rho : \mathcal{P} \times_G W := [\mathcal{P} \times W/G] \rightarrow \mathcal{C}$$

and the *universal section* $u : W \rightarrow \mathcal{P} \times_G W$ of ρ . This is summarized in the diagram:

$$\begin{array}{ccccccc} \mathcal{P} \times_G W & \xleftarrow[u]{\rho} & \mathcal{C} & \xrightarrow{\pi} & Q_{g,k}^\varepsilon(X, \beta) & \xrightarrow{\sigma} & \mathfrak{Bun}_G^{\text{tw}} \\ & & \downarrow [u] & & & & \\ & & [W/G] & & & & \end{array}$$

Let \mathbb{T}_ρ be the relative tangent complex of ρ . The σ -relative obstruction theory is given by the complex $(R^\bullet \pi_* u^* \mathbb{T}_\rho)^\vee$. As shown in Theorem 4.5.2 of [12], this complex is two-term perfect if W is LCI and W^s is smooth.

From now on we will assume that W is LCI and $W^s = W^{ss}$ is smooth.

Note that there are quasi-isomorphisms

$$R^\bullet \pi_* (\mathcal{P} \times_G \mathfrak{g} \cong u^*(\mathcal{P} \times_G (\mathfrak{g} \otimes \mathcal{O}_W))) \simeq (\sigma^* \mathbb{L}_{\mathfrak{Bun}_G^{\text{tw}}/\mathfrak{M}_{g,k}^{\text{tw}}}[1])^\vee$$

and $\mathbb{T}_\rho \simeq \mathcal{P} \times_G \mathbb{T}_W$.

Note also that on W there is a natural distinguished triangle

$$\mathfrak{g} \otimes \mathcal{O}_W \rightarrow \mathbb{T}_W \rightarrow \mathbb{T}_{[W/G]|_W}.$$

Therefore, we obtain the perfect obstruction theory

$$(R^\bullet \pi_* [u]^* \mathbb{T}_{[W/G]})^\vee$$

for $Q_{g,k}^\varepsilon(X, \beta)$ relative to the pure dimensional, smooth stack $\mathfrak{M}_{g,k}^{\text{tw}}$ as in [8, §5.3]. Further, the two relative perfect obstruction theories determine the same absolute perfect obstruction theory on $Q_{g,k}^\varepsilon(X, \beta)$ and the virtual classes associated to all three obstruction theories coincide.

2.5. Basic Properties and Variants.

2.5.1. *Expected dimension.* Since the étale gerbe markings are away from base locus for ε -stable quasimaps, there are natural evaluation morphisms

$$ev_i : Q_{g,k}^\varepsilon(X, \beta) \rightarrow \bar{I}_\mu X, ((C, x_1, \dots, x_k), [u]) \mapsto [u]_{|x_i} \quad i = 1, \dots, k,$$

to the rigidified cyclotomic inertia stack $\bar{I}_\mu X$ of the DM-stack X , as explained in [1, §4.4]. The stack $\bar{I}_\mu X$, which parameterizes representable maps from gerbes banded by finite cyclic groups to X , is a smooth (resp. proper) stack over \mathbb{C} if X is a smooth (resp. proper) stack over \mathbb{C} (see [1, Corollary 3.4.2]).

Let $\coprod_{c \in R} X_c$ denote the connected component decomposition of $\bar{I}_\mu X$ for some index set R and, for $c_i \in R, i = 1, \dots, k$, let

$$Q_{g,k}^\varepsilon(X, \beta; c_1, \dots, c_k) := \left(\prod_i ev_i \right)^{-1} \left(\prod_i X_{c_i} \right).$$

Its virtual dimension is

$$(2.5.1) \quad k + (1 - g)(\dim X - 3) + \beta(\det \mathbb{T}_{[W/G]}) - \sum_{i=1}^k \text{age} X_{c_i}$$

by Riemann-Roch for twisted curves ([1, Theorem 7.2.1]). The age is defined as follows. Let (\bar{x}, g) be a geometric point of X_c , with r the order of $g \in \text{Aut}(\bar{x})$. The age of X_c is

$$\sum_{j=0}^{r-1} \frac{j}{r} \dim E_j$$

where E_j is the eigenspace of $T_{\bar{x}} X$ of the induced μ_r -action with eigenvalue $e^{2\pi\sqrt{-1}j/r}$.

2.5.2. *Trivializations of gerbe markings.* As in [1, §6.1.3], one may construct the moduli stacks whose objects over a scheme S are S -families of ε -stable quasimaps *with sections of the gerbe markings*. We will not use these stacks.

2.5.3. *Graph spaces.* For $g, k \geq 0$ and $\varepsilon \geq 0+$, define the graph space $QG_{g,k,\beta}^\varepsilon(X)$ to be the moduli stack of ε -stable graph quasimaps. By a (resp. prestable) graph quasimap we mean the data

$$((C, x_1, \dots, x_k), P, [u] := ([u]_1, [u]_2))$$

with $((C, x_1, \dots, x_k), [u]_1)$ a k -pointed, genus g (resp. prestable) quasimap to X and a map $[u]_2 : C \rightarrow \mathbb{P}^1$ for which $[u]_2 : \underline{C} \rightarrow \mathbb{P}^1$ is a degree 1 map, that is, there is a unique component of \underline{C} isomorphic to \mathbb{P}^1 under $[u]_2$. For $\varepsilon \in \mathbb{Q}_{>0}$, the ε -stability for a prestable graph quasimap is defined by imposing the requirements that

$$(2.5.2) \quad \omega_{\underline{C}}(\sum x_i) \otimes (\phi_*([u]_1^* L_\theta)^e)^{\varepsilon/e} \otimes [u]_2^* \mathcal{O}_{\mathbb{P}^1}(3) \text{ is ample}$$

and that

$$(2.5.3) \quad \varepsilon l(x) \leq 1, \text{ for every } \mathbb{C}\text{-point } x \in C.$$

Again, $l(x)$ in (2.5.3) is the length of the quasimap at x defined in [12, §7.1]. When the requirement (2.5.2) is true for every small enough $\varepsilon \in \mathbb{Q}_{>0}$, we say that the graph quasimap is $(0+)$ -stable (the length inequality imposes no condition and is discarded for $\varepsilon = 0+$).

For $\varepsilon \in [0+, \infty)$, by the same argument as in §2.4, $QG_{0,k,\beta}^\varepsilon(X)$ is a finite type, proper DM-stack over \underline{X}_0 . The universal family of ε -stable graph quasimaps consists of the universal principal bundle \mathcal{P} on the universal curve $\pi : \mathcal{C} \rightarrow QG_{g,k,\beta}^\varepsilon(X)$ and the universal map

$$u = (u_1, [u_2]) : \mathcal{C} \rightarrow (\mathcal{P} \times_G W) \times \mathbb{P}^1,$$

with u_1 the universal section of $\rho : \mathcal{P} \times_G W \rightarrow \mathcal{C}$. If we let

$$\tilde{\rho} : (\mathcal{P} \times_G W) \times \mathbb{P}^1 \rightarrow \mathcal{C}$$

be the composition of ρ with the projection to the first factor, then the perfect obstruction theory relative to $\mathfrak{Bun}_G^{\text{tw}}$ remains of the same form $(R^\bullet \pi_* u^* \mathbb{T}_{\tilde{\rho}})^\vee$.

2.5.4. *Localization.* Assume W has an action by an algebraic torus T , commuting with the G -action. This induces T -actions on \mathcal{X} , X , \underline{X}_0 , $\bar{I}_\mu X$, and on the moduli spaces $Q_{g,k}^\varepsilon(X, \beta)$. In this situation, there is a T -equivariant embedding of $Q_{g,k}^\varepsilon(X, \beta)$ into a smooth Deligne-Mumford stack. The argument from §6.3 of [12] works. Therefore we may apply the virtual localization theorem of [17] to $Q_{g,k}^\varepsilon(X, \beta)$ and $QG_{g,k,\beta}^\varepsilon(X)$.

2.5.5. *Orbifold quasimaps with weighted markings.* The results of §2.3, §2.4 also provide the extension to orbifold targets of the theory of quasimaps with weighted markings developed in [11].

Let θ_0 be the minimal *integral* character on the ray $\theta \cdot \mathbb{Q}_{>0} \subset \chi(G)_{\mathbb{Q}}$ in the character group of G with \mathbb{Q} -coefficients. Then the ε -stability condition on quasimaps can be interpreted as stability with respect to the *rational* character $\varepsilon\theta_0$, see Definition 2.6 and Proposition 2.7 (i) in [11].

Now for $\varepsilon \in \mathbb{Q}_{>0} \cup \{0+\}$ and $\delta_1, \dots, \delta_m \in (\mathbb{Q}_{>0} \cap (0, 1]) \cup \{0+\}$, consider the rational character

$$\theta := \varepsilon\theta_0 + \sum \delta_j \text{id}_{\mathbb{C}^*}$$

of the group $G \times (\mathbb{C}^*)^m$. Then the moduli space of genus g , ε -stable quasimaps of class β to X , with k usual markings and m markings weighted by $\delta_1, \dots, \delta_m$, is identified with the moduli stack

$$Q_{g,k}^{\theta}([W^{ss}/G] \times [\mathbb{C}^{ss}/\mathbb{C}^*]^m, (\beta, 1, \dots, 1))$$

constructed in §2.3. Note that the domain curves carry nontrivial stack structure only at nodes and at the usual markings.

An important special case considered in [11] is when $\delta_1 = \dots = \delta_m = \delta$ for some $\delta \in (\mathbb{Q}_{>0} \cap (0, 1]) \cup \{0+\}$. This gives a theory with *two* stability parameters (ε, δ) , for which the corresponding moduli stacks are denoted $Q_{g,k|m}^{\varepsilon,\delta}(X, \beta)$.

Similarly, there are graph spaces with weighted markings.

By Theorem 2.3.7, the moduli spaces with weighted markings are DM stacks, proper over the affine quotient, and carrying the canonical perfect obstruction theory for LCI W .

3. QUASIMAP THEORY

3.1. **Quasimap Invariants.** We extend the quasimap theory for the orbifold case ($\mathfrak{X} = [W/G], X = [W^{ss}/G]$), closely following [1, 12, 9]. Fix an algebraic torus T action on W , commuting with the given G action. We allow the case when T is the trivial group. Assume that the T fixed locus \underline{X}_0^T of the affine quotient \underline{X}_0 is a finite set of points. Let $K := \mathbb{Q}(\{\lambda_i\})$ be the rational localized T -equivariant cohomology of $\text{Spec}\mathbb{C}$, with $\{\lambda_1, \dots, \lambda_{\text{rank}(T)}\}$ corresponding to a basis for the characters of T . The Novikov ring is defined to be

$$\Lambda_K := K[[\text{Eff}(W, G, \theta)]].$$

We write q^β for the element corresponding to β in Λ_K so that Λ_K is the q -adic completion. We denote by $q\Lambda_K$ the maximal ideal generated by q^β , $\beta \neq 0$.

Let $\{\gamma_i\}$ be a basis of the T -equivariant *Chen-Ruan cohomology* of X ,

$$H_{\text{CR},T}^*(X, \mathbb{Q}) := H_T^*(\bar{I}_\mu X, \mathbb{Q})$$

and let $\{\gamma^i\}$ be the dual basis “with respect to the Poincaré pairing in the *non-rigidified* cyclotomic inertia stack $I_\mu X$ of X ” in the sense that

$$\langle \gamma_i, \gamma^j \rangle_{\text{orb}} := \int_{\sum_{r \in \mathbb{N}_{\geq 1}} r^{-1} [\bar{I}_{\mu_r} X]} \gamma_i \cdot \iota^* \gamma^j = \delta_i^j,$$

with ι the involution of $\bar{I}_\mu X$ obtained from the inversion automorphisms. Note that

$$\sum_{r=1}^{\infty} r [\Delta_{\bar{I}_{\mu_r} X}] = \sum_i \gamma_i \otimes \gamma^i \text{ in } H^*(\bar{I}_\mu X \times \bar{I}_\mu X, \mathbb{Q}),$$

where the diagonal class $[\Delta_{\bar{I}_{\mu_r} X}]$ is obtained via push-forward of the fundamental class by $(\text{id}, \iota) : \bar{I}_{\mu_r} X \rightarrow \bar{I}_{\mu_r} X \times \bar{I}_{\mu_r} X$.

Define ψ_i to be the first Chern class of the universal cotangent line whose fiber at $((C, x_1, \dots, x_k), [u])$ is the cotangent space of the coarse moduli \underline{C} of C at i -th marking \underline{x}_i . For non-negative integers a_i and classes $\alpha_i \in H_T^*(\bar{I}_\mu X, \mathbb{Q})$, $t = \sum_j t_j \gamma_j$ with formal variables t_j , we write

$$\begin{aligned} \langle \alpha_1 \psi^{a_1}, \dots, \alpha_k \psi^{a_k} \rangle_{g,k,\beta}^\varepsilon &:= \int_{[Q_{g,k}^\varepsilon(X,\beta)]^{\text{vir}}} \prod_i ev_i^*(\alpha_i) \psi_i^{a_i}; \\ \langle\langle \alpha_1 \psi^{a_1}, \dots, \alpha_k \psi^{a_k} \rangle\rangle_{k,\beta}^\varepsilon &:= \sum_{m \geq 0} \frac{1}{m!} \langle \alpha_1 \psi^{a_1}, \dots, \alpha_k \psi^{a_k}, t, \dots, t \rangle_{g,k+m,\beta}^\varepsilon; \\ \langle\langle \alpha_1 \psi^{a_1}, \dots, \alpha_k \psi^{a_k} \rangle\rangle_k^\varepsilon &:= \sum_{\beta, m} \frac{q^\beta}{m!} \langle \alpha_1 \psi^{a_1}, \dots, \alpha_k \psi^{a_k}, t, \dots, t \rangle_{g,k+m,\beta}^\varepsilon \\ &\in \Lambda_K[[\{t_j\}_j]]. \end{aligned}$$

We may also define quasimap Chen-Ruan classes. Write

$$(3.1.1) \quad (\tilde{e}v_j)_* = \iota_*(\mathbf{r}_j(ev_j)_*),$$

where \mathbf{r}_j is the order function of the band of the gerbe structure at the marking j . Define a class in $H_*^T(\bar{I}_\mu X) \cong H_T^*(\bar{I}_\mu X)$ by

$$\begin{aligned} \langle \alpha_1, \dots, \alpha_k, - \rangle_{g,\beta}^\varepsilon &:= (\tilde{e}v_{k+1})_* \left(\left(\prod ev_i^* \alpha_i \right) \cap [Q_{g,k}^\varepsilon(X, \beta)]^{\text{vir}} \right) \\ &= \sum_i \gamma^i \langle \alpha_1, \dots, \alpha_k, \gamma_i \rangle_{g,k+1,\beta}^\varepsilon. \end{aligned}$$

Since the evaluation maps are proper, these are well-defined without T -localization, even when the coarse moduli space \underline{X} is not projective.

3.2. J^ε -function. As in [8, 9], we use \mathbb{C}^* -residues on graph spaces to define J^ε -functions. However, simply copying [9, Definition 5.1.1] in the orbifold setting would give a function with values in the untwisted sector $H_T^*(X)$. To ensure that the J^ε -functions take values in the full Chen-Ruan cohomology of X , including the twisted sectors, we will use instead graph spaces with an extra marking.

First of all, we fix a \mathbb{C}^* action on \mathbb{P}^1 given by

$$t[\zeta_0, \zeta_1] = [t\zeta_0, \zeta_1],$$

for $t \in \mathbb{C}^*$, $[\zeta_0, \zeta_1] \in \mathbb{P}^1$. We consider $QG_{g,k,\beta}^\varepsilon(X)$ with the induced \mathbb{C}^* action as well as the induced T action. This gives rise to the canonical $T \times \mathbb{C}^*$ -equivariant perfect obstruction theory on $QG_{g,k,\beta}^\varepsilon(X)$ via the action of $T \times \mathbb{C}^*$ on the tangent complex $\mathbb{T}_{\mathfrak{X}} \boxtimes T_{\mathbb{P}^1}$.

We use the same notation for the evaluation maps on graph spaces

$$ev_i : QG_{g,k,\beta}^\varepsilon(X) \rightarrow \bar{I}_\mu X \times \mathbb{P}^1, \quad i = 1, \dots, k.$$

Let z denote the \mathbb{C}^* -equivariant Euler class of the line bundle associated to the fundamental representation space of \mathbb{C}^* .

We define $T \times \mathbb{C}^*$ -equivariant invariants on the graph spaces as follows: for $\sigma_j \in H_T^*(\bar{I}_\mu X) \otimes H_{\mathbb{C}^*}^*(\mathbb{P}^1)$, $t \in H_T^*(\bar{I}_\mu X) \cong H_T^*(\bar{I}_\mu X) \otimes 1 \subset H_T^*(\bar{I}_\mu X) \otimes H_{\mathbb{C}^*}^*(\mathbb{P}^1)$

$$\begin{aligned} \langle \sigma_1, \dots, \sigma_k \rangle_{k,\beta}^{QG^\varepsilon} &:= \int_{[QG_{0,k,\beta}^\varepsilon(X)]^{\text{vir}}} \prod_i ev_i^*(\sigma_i); \\ \langle\langle \sigma_1, \dots, \sigma_k \rangle\rangle_{k,\beta}^{QG^\varepsilon} &:= \sum_{m \geq 0} \frac{1}{m!} \langle \sigma_1, \dots, \sigma_k, t, \dots, t \rangle_{k+m,\beta}^{QG^\varepsilon}; \\ \langle\langle \sigma_1, \dots, \sigma_k \rangle\rangle_k^{QG^\varepsilon} &:= \sum_{\beta, m} \frac{q^\beta}{m!} \langle \sigma_1, \dots, \sigma_k, t, \dots, t \rangle_{k+m,\beta}^{QG^\varepsilon}. \end{aligned}$$

The integration is understood as equivariant push-forward to $\text{Spec} \mathbb{C}$, hence $\langle \sigma_1, \dots, \sigma_k \rangle_{k,\beta}^{QG^\varepsilon} \in K[z]$ and the double brackets are elements of $K[[z]][[\text{Eff}(W, G, \theta)]][[\{t_i\}_i]]$.

In what follows, for an integer $k \geq 0$, $[k]$ will mean the index set $\{1, \dots, k\}$. Let $F := QG_{0,[k] \cup \{\star\}, \beta}^\varepsilon(X)^{\mathbb{C}^*}$ denote the \mathbb{C}^* -fixed substack of $QG_{0,[k] \cup \{\star\}, \beta}^\varepsilon(X)$ (the maximal closed substack fixed by the action of every element of \mathbb{C}^*). If $(C, x_\star, \mathbf{x}, [u])$ is a \mathbb{C}^* -fixed ε -stable graph quasimap, then on the coarse curve of the distinguished component of C we must have that $[u]_{\text{reg}}$ is a constant map to \underline{X} . The base locus of $[u]$, if nonempty, must be over $0 := [0, 1]$ or $\infty := [1, 0]$ in the target \mathbb{P}^1 . The same is true for the markings (x_\star, \mathbf{x}) .

As in [9, §4.2], there are “components” $F_{A_2, \beta_2}^{A_1, \beta_1}$ of F corresponding to distributions of the markings and of the class β over 0 and ∞ : for $A_1 \coprod A_2 = [k] \cup \{\star\}$ and $\beta_1 + \beta_2 = \beta$, the closed substack $F_{A_2, \beta_2}^{A_1, \beta_1}$ parameterizes \mathbb{C}^* -fixed ε -stable graph quasimaps for which markings in A_1 and the class β_1 are supported over 0, while A_2, β_2 are supported over ∞ . In particular, we have a distinguished \mathbb{C}^* -fixed part $F_{\star, 0}^{k, \beta}$ of $QG_{0, [k] \cup \{\star\}, \beta}^\varepsilon(X)$. This closed substack $F_{\star, 0}^{k, \beta}$ consists of ε -stable graph quasimaps $((C, x_\star, \mathbf{x}), [u])$ such that only the marking x_\star is over ∞ while the markings in $[k]$ and the entire class β are over $0 \in \mathbb{P}^1$.

Let η_0, η_∞ be the \mathbb{C}^* -equivariant classes of \mathbb{P}^1 defined by the property

$$\eta_0|_0 = z, \quad \eta_\infty|_\infty = -z, \quad \eta_0|_\infty = 0 = \eta_\infty|_0.$$

Definition 3.2.1. 1. Define the J^ε -function by

$$J^\varepsilon(t, q, z) := \sum_{\beta \in \text{Eff}(W, G, \theta), k \geq 0} q^\beta (\text{pr}_{\bar{I}_\mu X} \widetilde{\circ} ev_\star)_* \left(ev_\star^*(\eta_\infty) \frac{\prod_{i=1}^k ev_i^*(t)}{k!} \frac{[F_{\star, 0}^{k, \beta}]^{\text{vir}}}{e^{\mathbb{C}^* \times T}(N_{F_{\star, 0}^{k, \beta} / QG_{0, [k] \cup \{\star\}, \beta}^\varepsilon(X)})} \right),$$

where pr_X denotes the projection to X from $X \times \mathbb{P}^1$ and the notation $(\text{pr}_{\bar{I}_\mu X} \widetilde{\circ} ev_\star)_*$ is as defined in (3.1.1). The localization residue is taken as a sum over the connected components of $F_{\star, 0}^{k, \beta}$.

2. Define the S^ε -operator by

$$\begin{aligned} S_t^\varepsilon(z)(\gamma) &:= \sum_i \gamma_i \left\langle \frac{\gamma^i}{z - \psi}, \gamma \right\rangle_{0, \{\bullet, \star\}}^\varepsilon \\ &:= \gamma + \sum_{\beta \neq 0 \text{ or } k \geq 1} q^\beta (\tilde{ev}_\bullet)_* \left(\frac{ev_\star^*(\gamma) \prod_{i=1}^k ev_i^*(t)}{z - \psi_\bullet} \frac{\prod_{i=1}^k ev_i^*(t)}{k!} \cap [Q_{0, [k] \cup \{\bullet, \star\}}^\varepsilon(X, \beta)]^{\text{vir}} \right) \\ &= \gamma + O(1/z). \end{aligned}$$

Remark 3.2.2. (1) It is straightforward to check that the definitions of J^ε and S^ε agree with those given for the same objects in [8, 9] when X becomes a scheme.

- (2) As noted before, properness of the evaluation maps implies that J^ε and S^ε are well-defined without T -localization for all targets.
- (3) When $\beta = 0$ the moduli spaces coincide for different values of ε coincide. Hence $J^\varepsilon(t, q, z)$, $S_t^\varepsilon(z)$ modulo q do not depend on ε .
- (4) Consider Givental’s symplectic space

$$\mathcal{H}_X := H_T^*(\bar{I}_\mu X) \otimes_{\mathbb{Q}} K((z^{-1}))[[\text{Eff}(W, G, \theta)]].$$

The series J^ε is an element of $\mathcal{H}_X[[\{t_j\}_j]]$ and the operator S_t^ε may be viewed as a family (parametrized by the formal variables t_j) of endomorphisms of \mathcal{H}_X .

3.3. Factorization. For an Artin stack \mathcal{Y} over \mathbb{C} , a twisted curve C (with some gerbe markings) over \mathbb{C} and $\beta \in \text{Hom}_{\mathbb{Z}}(\text{Pic}\mathcal{Y}, \mathbb{Q})$, we denote by $\text{Hom}_{\beta}^{\text{rep}}(C, \mathcal{Y})$ the substack of $\text{Hom}_{\text{Spec}\mathbb{C}}(C, \mathcal{Y})$ consisting of representable 1-morphisms $C \rightarrow \mathcal{Y}$ with class β .

Let μ_r be the group of r -th roots of unity. We will consider a special twisted curve

$$C_{r,-r} := [\mathbb{P}^1/\mu_r]$$

where μ_r acts \mathbb{P}^1 by $s \cdot [\zeta_0, \zeta_1] = [s\zeta_0, \zeta_1]$ for $s \in \mu_r$, $[\zeta_0, \zeta_1] \in \mathbb{P}^1$.

Lemma 3.3.1. *Let Y be a DM stack over \mathbb{C} .*

- (1) *Consider a representable morphism $[u] := ([u]_1, [u]_2)$ from a genus 0 irreducible twisted curve (C, x_1, x_2) over a scheme S to $Y \times \mathbb{P}^1$ sending x_1 to 0 and x_2 to ∞ under $[u]_2$, with class $(0, [\mathbb{P}^1])$. Then for some positive integer r , C is canonically isomorphic to $C_{r,-r} \times S$ with $[u]_2$ as the coarse moduli space.*
- (2) *The closed substack $(\text{pr}_{\mathbb{P}^1} \circ \text{ev}_1)^{-1}(0) \cap (\text{pr}_{\mathbb{P}^1} \circ \text{ev}_2)^{-1}(\infty)$ of $\mathcal{K}_{0,2}(Y \times \mathbb{P}^1, (0, [\mathbb{P}^1]))$ is isomorphic to the Hom stack*

$$\text{Hom}_0^{\text{rep}}(C_{r,-r}, Y)$$

parameterizing representable 1-morphisms from $C_{r,-r}$ to Y with class 0.

- (3) *The stack $\text{Hom}_0^{\text{rep}}(C_{r,-r}, Y)$ is isomorphic to the cyclotomic inertia stack $I_\mu Y$ of Y , compatible with the rigidification map $I_\mu Y \rightarrow \bar{I}_\mu Y$ and the two evaluation maps (with the involution). Here $\text{pr}_{\mathbb{P}^1}$ denotes the projection to \mathbb{P}^1 from $Y \times \mathbb{P}^1$.*

Proof. For 1): By [1, Theorem 4.2.1], it is enough to show the case when $S = \text{Spec}\mathbb{C}$. We may assume that $Y = B\Gamma$ for some finite group Γ . Denote by \bar{x}_i the geometric points associated to the gerbes x_i . The monodromies of the induced Γ -bundle on an étale chart of C around \bar{x}_i are exactly the homomorphisms $\text{Aut}(\bar{x}_i) \rightarrow \text{Aut}(\bar{Y}) = \Gamma$, where \bar{Y} is the \mathbb{C} -point of Y . Since the homomorphisms are monomorphisms, we are done.

For 2): There is a natural functor Φ from $\text{Hom}_0^{\text{rep}}(C_{r,-r}, Y)$ to $\mathcal{D} := (\text{pr}_{\mathbb{P}^1} \circ \text{ev}_1)^{-1}(0) \cap (\text{pr}_{\mathbb{P}^1} \circ \text{ev}_2)^{-1}(\infty)$. It is essentially surjective by (1). It is clearly faithful. Given an arrow (φ, α) in \mathcal{D} between objects $(C_{r,-r} \times S, [u])$ and $(C_{r,-r} \times S, [u'])$ over a scheme S , i.e., an 1-isomorphism $\varphi : C_{r,-r} \times S \rightarrow C_{r,-r} \times S$ over S with a 2-morphism $\alpha : [u] \Rightarrow [u'] \circ \varphi$, first note that there is a unique 2-morphism $\sigma : \varphi \Rightarrow \text{id}_{C_{r,-r} \times S}$. Now

define $\alpha' := [u'](\sigma) \circ \alpha$ so that $\sigma : (\varphi, \alpha) \Rightarrow (\text{id}_{C_{r,-r} \times S}, \alpha')$. This shows that the functor Φ is full.

For 3): Every object $[\mathbb{P}^1/\mu_r] \times S \rightarrow Y$ in $\text{Hom}_0^{\text{rep}}(C_{r,-r}, Y)$ is factored through a unique representable morphism $B\mu_r \times S \rightarrow Y$ given by one of sections of the inclusions $x_1, x_2 \subset [\mathbb{P}^1/\mu_r] \times S$. This yields a functor $\text{Hom}_0^{\text{rep}}(C_{r,-r}, Y) \rightarrow I_\mu Y$. In fact, this functor is an equivalence according to the definition [1, Definition 3.2.1] of $I_\mu Y$. \square

Our next goal is to prove the following factorization of graph quasimap double brackets.

Proposition 3.3.2. *Let $\gamma, \sigma \in H_T^*(\bar{I}_\mu X)$. Then*

(3.3.1)

$$\langle\langle \gamma \otimes \eta_0, \delta \otimes \eta_\infty \rangle\rangle_{\{\star_1, \star_2\}}^{QG^\varepsilon} = \sum \langle\langle \gamma, \frac{\gamma^i}{z - \psi} \rangle\rangle_{0, \{\star_1, \bullet\}}^\varepsilon \langle\langle \frac{\gamma^i}{-z - \psi}, \delta \rangle\rangle_{0, \{\bullet, \star_2\}}^\varepsilon$$

We will use \mathbb{C}^* -localization, as in the proof Proposition 5.3.1 of [9]. To this end, we first establish a factorization of the virtual classes of the \mathbb{C}^* -fixed loci $F_{A_2, \beta_2}^{A_1, \beta_1}$.

We use 0 and ∞ to label the two markings \star_1, \star_2 by abusing notation. Let $D^{\text{tw}}(A_1, A_2)$ denote the locally finite type smooth Artin stack parameterizing genus 0 nodal twisted curves with two *distinguished nodes*, splitting the curve into three twisted curves $(C_1, A_1 \cup \{\bullet\})$, $(C_{r, -r}, \{\bullet, \check{\bullet}\})$, and $(C_2, A_2 \cup \{\check{\bullet}\})$, together with sections of the gerbes $\bullet, \check{\bullet}$ (see [1, §5.1]). Here $(C_{r, -r}, \{\bullet, \check{\bullet}\})$ denotes $C_{r, -r}$ for r =the index of the band of the gerbe \bullet .

Let $A_1 \amalg A_2 = [k] \cup \{0, \infty\}$. Assume $0 \in A_1, \infty \in A_2$. Below, the unstable terms $Q_{0, \{0, \bullet\}, \beta_1=0}^\varepsilon(X)$, $Q_{0, \{\infty, \check{\bullet}\}, \beta_2=0}^\varepsilon(X)$ are understood as $\bar{I}_\mu X$.

Consider the commuting diagram:

$$\begin{array}{ccccc} Q(A_1, A_2, \beta) & \xrightarrow[\cong]{\Psi_1} & F_{A_1, A_2, \beta} & \longrightarrow & QG_\beta^\varepsilon \\ \downarrow & & \downarrow & & \downarrow \\ D^{\text{tw}}(A_1, A_2) & \xrightarrow[\cong]{\Psi_2} & \mathfrak{M}_{A_1, A_2}^{\text{tw}, \mathbb{C}^*} & \longrightarrow & \mathfrak{M}_{0, k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1]), \end{array}$$

where

$$\begin{aligned} Q(A_1, A_2, \beta) &:= \coprod_{\beta_1 + \beta_2 = \beta} Q_{0, A_1 \cup \{\bullet\}, \beta_1}^\varepsilon(X) \times_{\bar{I}_\mu X} I_\mu X \times_{\bar{I}_\mu X} Q_{0, A_2 \cup \{\check{\bullet}\}, \beta_2}^\varepsilon(X); \\ F_{A_1, A_2, \beta} &:= \coprod_{\beta_1 + \beta_2 = \beta} F_{A_2, \beta_2}^{A_1, \beta_1}; \\ QG_\beta^\varepsilon &:= QG_{0, k, \beta}^\varepsilon(X). \end{aligned}$$

Further,

$$\mathfrak{M}_{0,k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1]) \subset \text{Hom}_{\mathfrak{M}_{0,k}^{\text{tw}}}(\mathfrak{C}, \mathbb{P}^1 \times \mathfrak{M}_{0,k}^{\text{tw}})$$

is the (locally finite type and smooth) Artin stack of k -pointed, genus 0, twisted (not necessarily stable) maps to \mathbb{P}^1 with class $[\mathbb{P}^1]$, while $\mathfrak{M}_{A_1, A_2}^{\text{tw}, \mathbb{C}^*}$ is the part of the \mathbb{C}^* -fixed closed substack of $\mathfrak{M}_{0,k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1])$ for which the markings in A_1 are over 0 and the markings in A_2 are over ∞ .

The vertical arrows are the natural morphisms obtained by forgetting the mapping data to \mathfrak{X} .

The two right horizontal arrows are the inclusions of the maximal \mathbb{C}^* -fixed closed substacks into their ambient stacks.

The two left horizontal arrows Ψ_1, Ψ_2 are obtained from gluing morphisms, constructed as follows. By Lemma 3.3.1, an object in $Q(A_1, A_2, \beta)$ can be considered as a triple

$$((C_1, \mathbf{x}_{A_1}, x_\bullet), [u]_1), ((C_0, x_\bullet, x_\bullet), [u]_0), ((C_2, x_\bullet, \mathbf{x}_{A_2}), [u]_2)$$

of ε -stable quasimaps to X , ε -stable graph quasimaps to X , ε -stable quasimaps to X , respectively, with isomorphisms $x_\bullet \cong x_\bullet$ inverting the band structures (for each “join”). With these isomorphisms we glue C_1, C_0, C_2 along x_\bullet, x_\bullet to get a twisted curve over S :

$$C_1 \underset{x_\bullet}{\amalg} C_0 \underset{x_\bullet}{\amalg} C_2$$

(see [1, §A] for gluing of Artin stacks along closed substacks). By the co-cartesian property the mapping data $[u]_1 : C_1 \rightarrow \mathfrak{X}$ (followed by the inclusion $\mathfrak{X} \cong \mathfrak{X} \times [0, 1] \subset \mathfrak{X} \times \mathbb{P}^1$), $[u]_0 : C_0 \rightarrow X \times \mathbb{P}^1$, $[u]_2 : C_2 \rightarrow \mathfrak{X}$ (followed by the inclusion $\mathfrak{X} \cong \mathfrak{X} \times [1, 0] \subset \mathfrak{X} \times \mathbb{P}^1$) can be glued. This explains Ψ_1 . The construction of Ψ_2 is similar. By investigating the formal deformation spaces we note that Ψ_2 is an 1-isomorphism. So is Ψ_1 , since the left square in the diagram is cartesian.

Proposition 3.3.3. *Let*

$$\Delta_2 : \bar{I}_\mu X \times \bar{I}_\mu X \rightarrow (\bar{I}_\mu X \times \bar{I}_\mu X) \times (\bar{I}_\mu X \times \bar{I}_\mu X)$$

be the product of the diagonal map $(\text{id}, \iota) : \bar{I}_\mu X \rightarrow \bar{I}_\mu X \times \bar{I}_\mu X$. Then

$$\begin{aligned} [F_{A_1, A_2, \beta}]^{\text{vir}} &= \sum_{A_1 \amalg A_2 = [k], \beta_1 + \beta_2 = \beta} \\ &\Delta_2^!([Q_{0, A_1 \cup \{\bullet\}}^\varepsilon(X, \beta_1)]^{\text{vir}} \times [I_\mu X] \times [Q_{0, A_2 \cup \{\bullet\}}^\varepsilon(X, \beta_2)]^{\text{vir}}). \end{aligned}$$

Proof. First note that $\mathfrak{M}_{0,k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1])$ is smooth over $\mathfrak{M}_{0,k}^{\text{tw}}$ so that there is a canonical quasi-isomorphism

$$(\mathbb{L}_{\mathfrak{M}_{0,k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1])} \big|_{\mathfrak{M}_{A_1, A_2}^{\text{tw}, \mathbb{C}^*}})^{\mathbb{C}^*\text{-fixed}} \cong \mathbb{L}_{\mathfrak{M}_{A_1, A_2}^{\text{tw}, \mathbb{C}^*}}.$$

Therefore, if we denote by E the absolute perfect obstruction theory for QG_β^ε , then the \mathbb{C}^* -fixed part of the distinguished triangle

$$E|_{F_{A_1, A_2, \beta}} \rightarrow (R^\bullet \pi_* [u_1]^* \mathbb{T}_{\mathfrak{X}})^\vee \rightarrow \mathbb{L}\mathfrak{M}_{0,k}^{\text{tw}}(\mathbb{P}^1, [\mathbb{P}^1])|_{F_{A_1, A_2, \beta}}[1]$$

yields the compatible absolute and relative perfect obstruction theories. By the normalization sequence of nodal curves with distinguished nodes and the tangent bundle lemma [1, Lemma 3.6.1], we can apply the functoriality from [4, Proposition 7.5] to complete the proof. \square

Proof of Proposition 3.3.2. We compute the Euler class of the virtual normal bundle of $F_{A_1, A_2, \beta}$ in QG_β^ε when $0 \in A_1$, $\infty \in A_2$. Assume $\beta_i \neq 0$, $i = 1, 2$. Then the Euler class comes from two contributions:

- (1) the moving part of the deformation of $u_2 : C \rightarrow \mathbb{P}^1$;
- (2) the moving part of the deformation and the infinitesimal automorphism of (C, \mathbf{x}) .

Part (1) contributes z (respectively, contributes $-z$) for each connected component of $C \setminus C_0$ contracted to 0 (respectively, contracted to ∞) under $[u]_2$.

Part (2) has contributions exactly from $T_{\bar{p}_i} C_1 \otimes T_{\bar{p}_0, i} C_0$ for each separating node \bar{p}_i of C (here \bar{p}_i are the geometric points associated to the separating node). The contributions are $\frac{z-\psi_\bullet}{r}$ for \bar{p}_1 and $\frac{-z-\psi_\bullet}{r}$ for \bar{p}_2 if $r = |\text{Aut}(\bar{p}_i)|$.

When $\beta_1 = 0$, $|A_1| = 1$ and $\beta_2 \neq 0$ or $|A_2| > 1$ there is no contribution $\frac{z-\psi_\bullet}{r}$; this is exactly the case when the separating node \bar{p}_1 disappears. Similarly, when $\beta_2 = 0$, $|A_2| = 1$ and $\beta_1 \neq 0$ or $|A_1| > 1$, then the separating node \bar{p}_i , respectively $i = 1, 2$ disappears and there is no contribution $\frac{-z-\psi_\bullet}{r}$. Finally, when $\beta_1 = \beta_2 = 0$ and $|A_1| = |A_2| = 1$ both separating nodes disappear and there is neither contribution $\frac{z-\psi_\bullet}{r}$ nor $\frac{-z-\psi_\bullet}{r}$.

The factorization expression (3.3.1) is immediate by this analysis, Proposition 3.3.3, and the contributions of η_0 and η_∞ . \square

Lemma 3.3.4. *Fix $\varepsilon \in \mathbb{Q}_{>0}$. The J^ε -function of $X = [W^{ss}/G]$ takes the form*

$$\begin{aligned} J^\varepsilon(t, z) = & \mathbb{1}_X + \frac{t}{z} + \sum_{0 < \beta(L_\theta) \leq 1/\varepsilon} q^\beta (\text{pr}_{\bar{I}_\mu X} \widetilde{\circ} \text{ev}_\star)_* \frac{(-z)[F_{\star, 0}^{\theta, \beta}]^{\text{vir}}}{e^{\mathbb{C}^* \times T}(N_{F_{\star, 0}^{\theta, \beta}/QG_{0, \{\star\}, \beta}^\varepsilon(X)}^{\text{vir}})} \\ & + \sum_{\substack{(\beta \neq 0, k \geq 1) \text{ or} \\ (\beta(L_\theta) > 1/\varepsilon, k=0)}} q^\beta (\tilde{e}v_\bullet)_* \frac{\prod_{i=1}^k \text{ev}_i^*(t) \cap [Q_{0, [k] \cup \{\bullet\}}^\varepsilon(X, \beta)]^{\text{vir}}}{k! z(z - \psi_\bullet)}, \end{aligned}$$

where $\mathbb{1}_X$ is the fundamental class of X on the untwisted sector. Here again the localization residue is taken as a sum over the connected components of $F_{\star,0}^{\emptyset,\beta}$.

Proof. The first term is from the case $(k = 0, \beta = 0)$. In this case, $r = 1$ by Lemma 3.3.1 (1). Thus, $(F_{\star,0}^{0,0}, QG_{0,\star,0}^\varepsilon(X)) = (X \times \infty, X \times \mathbb{P}^1)$. This explains the first term.

The second term is from the case $(k = 1, \beta = 0)$. By Lemma 3.3.1 (2), $ev_\star : F_{\star,0}^{1,0} \rightarrow \bar{I}_\mu X$ is nothing but $\varpi : I_\mu X \rightarrow \bar{I}_\mu X$ under a suitable identification of $F_{\star,0}^{1,0} = I_\mu X$. Since the degree of $I_{\mu_r} X \rightarrow \bar{I}_{\mu_r} X$ is $1/r$, we obtain the second term.

The third sum is from the case $k = 0$ and $0 \neq \beta(L_\theta) \leq 1/\varepsilon$.

The last sum follows from the case when there is a separating node over $[0, 1] \in \mathbb{P}^1$, i.e., $k \geq 1$ or $\beta > 1/\varepsilon$ except $(k, \beta) = (1, 0)$. By Proposition 3.3.3 with one separating node over $[0, 1]$ and the analysis of the virtual normal bundle in the proof (3.3.1), the last sum is immediate. \square

Lemma 3.3.4 shows that J^∞ coincides with the J -function J defined in [28] after identifying $H^*(I_\mu X)$ and $H^*(\bar{I}_\mu X)$ by pullback (not by push-forward): $\varpi^*(zJ^\infty) = J$, where ϖ denotes the rigidification $I_\mu X \rightarrow \bar{I}_\mu X$.

3.4. Unitarity of S^ε . The following Proposition says that the operators S_t^ε are symplectic transformations on Givental's symplectic space \mathcal{H}_X .

Proposition 3.4.1. *Consider the operator*

$$(S^\varepsilon)_t^\star(-z)(\gamma) := \sum_i \gamma_i \langle\langle \gamma^i, \frac{\gamma}{-z - \psi} \rangle\rangle_{0, \{\star, \bullet\}}^\varepsilon = \gamma + O(1/z).$$

Then

$$(S^\varepsilon)^\star(-z) \circ S^\varepsilon(z)(\gamma) = \gamma.$$

Proof. The factorization (3.3.1), together with the fact that S^ε and $(S^\varepsilon)^\star$ are of form $\text{Id} + O(1/z)$, yields a proof as in [9, Proposition 5.3.1]. \square

3.5. The P -series and Birkhoff factorization of J^ε . Define the P^ε -series by

$$P^\varepsilon(t, q, z) := \sum \gamma_i \langle\langle \gamma^i \eta_\infty \rangle\rangle_{\{\star\}}^{QG^\varepsilon} = \mathbb{1}_X + O(q),$$

where the latter equality follows from Lemma 3.3.1.

Definition 3.5.1. We call the triple (W, G, θ) (or the “target” (\mathfrak{X}, X)) semi-positive if $\beta(\det T_{\mathfrak{X}})$ is non-negative for every $\beta \in \text{Eff}(W, G, \theta)$. (Note that $\beta(\det T_{[W/G]}) = \beta(\det T_W)$, since the determinant of adjoint bundle $P \times_G \text{Lie}(G)$ on the domain curve C has degree zero.)

Theorem 3.5.2. (1) The following formula holds:

$$J^\varepsilon(t, q, z) = S_t^\varepsilon(z)(P^\varepsilon(t, q, z)).$$

- (2) The small J -function $J^\varepsilon|_{t=0}$ has degree 0 if we set $\deg z = 1$, $\deg q^\beta = \beta(\det T_{\mathfrak{X}})$, and the degrees of cohomology classes of $\bar{I}_\mu X$ to be the age-shifted complex degrees.
- (3) For a semi-positive target (W, G, θ) , the following hold.
- (a) $J^\varepsilon(t, q, z)$ takes the form $J_0^\varepsilon(q)\mathbb{1}_X + \frac{1}{z}(t + J_1^\varepsilon(q)) + O(1/z^2)$, for some $J_0^\varepsilon(q) = 1 + O(q) \in \Lambda_K$ of degree 0 and $J_1^\varepsilon(q) \in (q\Lambda_K)H^{\leq 2}(\bar{I}_\mu X, \Lambda_K)$ of degree 1 with respect to the grading from (2) above.
- (b) $P^\varepsilon(t, q, z) = J_0^\varepsilon(q)\mathbb{1}_X$. In particular

$$\begin{aligned} \frac{J^\varepsilon(t, q, z)}{J_0^\varepsilon(q)} &= S_t^\varepsilon(\mathbb{1}_X) \\ \frac{t + J_1^\varepsilon(q)}{J_0^\varepsilon(q)} &= \sum \gamma_i \langle\langle \gamma^i, \mathbb{1}_X \rangle\rangle_{0,2}^\varepsilon - \mathbb{1}_X. \end{aligned}$$

- (c) $J_0^\varepsilon(q)\mathbb{1}_X$ is the unity in the ε -quasimap quantum product.
- (4) If $\beta(\det T_{\mathfrak{X}}) > 0$ for every nonzero L_θ -effective class, then $J^\varepsilon = \mathbb{1}_X + O(1/z)$.

Proof. The proof of (1) is completely analogous to the case when X is a nonsingular variety (see [9, Theorem 5.4.1]). Briefly, virtual localization of P^ε with respect to the \mathbb{C}^* action on graph spaces yields the factorization

$$(3.5.1) \quad P^\varepsilon = (S^\varepsilon)^*(-z)(J^\varepsilon).$$

Now apply $S^\varepsilon(z)$ to both sides of the above equality and use Proposition 3.4.1.

For (2), note that the virtual dimension of a connected component of $QG_{0,1,\beta}^\varepsilon$ landing on X_c under $\text{pr}_{\bar{I}_\mu X} \circ \text{ev}_*$ is, by the formula (2.5.1),

$$1 + (\dim \mathfrak{X} + 1 - 3) + \beta(\det T_{\mathfrak{X}}) + \deg[u]_2^* T_{\mathbb{P}}^1 - \text{age} X_c,$$

so that under $(\text{pr}_{\bar{I}_\mu X} \circ \tilde{e}v_*)^*$ after cap with η_∞ , its usual cohomological degree becomes $-\beta(\det T_{\mathfrak{X}}) - \dim \mathfrak{X} + \text{age} X_c + \dim X_c$. The age shifted degree of the latter becomes $-\beta(\det T_{\mathfrak{X}}) - \dim \mathfrak{X} + \text{age} X_c + \dim X_c + \text{age} \iota(X_c)$ which is $-\beta(\det T_{\mathfrak{X}})$.

For (3a), note that by (2) and Lemma 3.3.4, J^ε is a series of $1/z$. Considering the equality of (1) modulo $1/z$, we conclude that P^ε has only the zero-th power of z and coincides with the zero-th power of z piece in J^ε . The latter is $J_0^\varepsilon(q)\mathbb{1}_X$ by (2) for some $J_0^\varepsilon(q) \in \Lambda_K$.

Claim (3b) follows from (1) and (3a) (see [9, Corollary 5.5.3]). Claim (3c) follows from the identical proof of [9, Corollary 5.5.4].

Claim (4) is obvious from (2) and Lemma 3.3.4. \square

Remark 3.5.3. When $\varepsilon = \infty$, combining the relation (3.5.1) from the proof of Theorem 3.5.2 (1) with Lemma 3.3.4, we obtain

$$J^\infty = S^\infty(\mathbb{1}_X).$$

3.6. ε -wall-crossing. As generalizations of Theorems 7.3.1, 7.3.4 and Conjecture 6.2.1 of [9], it is natural to make the following conjecture.

Conjecture 3.6.1. (1) *The following formula holds:*

$$S_t^\varepsilon(\mathbb{1}_X) = S_{\tau(t)}^\infty(\mathbb{1}_X) \text{ for } \tau(t) := t + \sum_{\beta \neq 0} q^\beta \gamma_i \langle \langle \gamma^i, \mathbb{1}_X \rangle \rangle_{0, \{\bullet, \star\}, \beta}^\varepsilon.$$

(2) *The following formula holds:*

$$J^\varepsilon(t, q, z) = S_t^\varepsilon(z)(P^\varepsilon(t, q, z)) = S_{\tau^\infty, \varepsilon(t)}^\infty(z)(P^{\infty, \varepsilon}(\tau^{\infty, \varepsilon}(t), q, z))$$

for a unique transformation

$$t \mapsto \tau^{\infty, \varepsilon}(t) \in H_T^*(\bar{I}_\mu X) \otimes_{\mathbb{Q}} K[[\text{Eff}(W, G, \theta)]][[\{t_i\}_i]]$$

and a unique element

$$P^{\infty, \varepsilon}(t, q, z) \in H_T^*(\bar{I}_\mu X) \otimes_{\mathbb{Q}} K[z][[\text{Eff}(W, G, \theta)]][[\{t_i\}_i]]$$

In particular, zJ^ε is on the Lagrangian cone of the Gromov-Witten theory of X defined in [28] (with Novikov variables from Λ) and if the triple (W, G, θ) is semi-positive, then

$$J^\varepsilon(t, q, z)/J_0^\varepsilon(q) = J^\infty((t + J_1^\varepsilon(q))/J_0^\varepsilon(q), q, z)$$

with $\tau^{\infty, \varepsilon}(t) = (t + J_1^\varepsilon(q))/J_0^\varepsilon(q)$ and $P^{\infty, \varepsilon}(t, q, z) = J_0^\varepsilon(q)\mathbb{1}_X$.

The first main result of the paper is a proof of Conjecture 3.6.1 in the presence of a torus action with good properties.

Theorem 3.6.2. *Suppose that there is an action by an algebraic torus T on W which commutes with the G action and such that the induced T action on the coarse moduli space \underline{X} of X has only isolated fixed points. Then Conjecture 3.6.1 (1) holds true.*

Further, Conjecture 3.6.1 (2) holds true if we assume in addition that the 1-dimensional T -orbits are isolated when (W, G, θ) is not semi-positive.

3.7. Proof of Theorem 3.6.2. For X a scheme, the result is contained in [9, Theorems 7.3.1, 7.3.4] and the proof given there also works for orbifolds. We outline the argument, focusing on the appropriate changes.

3.7.1. *Unbroken quasimaps.* (cf. [9, §7.4.3]) For each \mathbb{C} -element in $Q_{0,2+m}^\varepsilon(X, \beta)^T$, we call the element is of *initial type* (resp. of *recursion type*) if the cotangent T -weight of the first marking at the coarse domain curve is zero (resp. nonzero). A recursion element is called *unbroken* if

$$\alpha_{C',x} + \alpha_{C'',x} = 0$$

for every node x of the domain curve C connecting C' and C'' , where $\alpha_{C',x}$, $\alpha_{C'',x}$ are the induced T -weights of the cotangent spaces at x to the coarse curves C' and C'' . A recursion element is called *broken* if there is a node x of the domain curve C connecting C' and C'' with $\alpha_{C',x} + \alpha_{C'',x} \neq 0$. Note that the domain curve of an unbroken recursion element has no components contracted under $[u]_{\text{reg}}$.

Let M be a connected component of $Q_{0,2}^\varepsilon(X, \beta)^T$. We call M a *recursion component* of $Q_{0,2}^\varepsilon(X, \beta)^T$ if it contains an unbroken element. This implies that every element in M is unbroken and is therefore a two-pointed *stable map to X* . Hence M is canonically identified with a connected component of $\mathcal{K}_{0,2}(X, \beta)^T$.

3.7.2. *Recursion.* (cf. [9, §7.5]) We divide the connected components of the T -fixed substack

$$Q_{0,2+m}^\varepsilon(X, \beta)^T$$

into initial types and recursion types, according to whether the first marking is on a contracted component or not under $[u]_{\text{reg}}$. Every recursion component is of the form $M \cong M' \times_{\bar{I}_\mu X^T} M''$, with M' an unbroken component of $\mathcal{K}_{0,2}(X, \beta')$ for some β' and M'' a connected component of $Q_{0,2+m}^\varepsilon(X, \beta - \beta')$, such that $\alpha_M = \alpha_{M'} \neq \alpha_{M''}$. Here α_M , $\alpha_{M'}$, $\alpha_{M''}$ denote the respective cotangent weights at the first coarse markings of any element in M , M' , M'' .

Let

$$\{\nu, \nu', \dots\} = \underline{\bar{I}_\mu X^T}$$

be the finite set of T -fixed \mathbb{C} -points of $\bar{I}_\mu X$ and let

$$i_\nu : \nu = \text{Spec } \mathbb{C} \rightarrow \bar{I}_\mu X$$

be the associated map. For $\gamma \in H_T^*(\bar{I}_\mu X) \otimes \Lambda_K$, denote

$$(3.7.1) \quad S_\nu^\varepsilon(\gamma) = i_\nu^*(S_t^\varepsilon(z)(\gamma)) \in K[[1/z]][[\text{Eff}(W, G, \theta)]][[t_j]].$$

Calculating $S_\nu^\varepsilon(\gamma)$ by virtual localization and using the above analysis of $Q_{0,2+m}^\varepsilon(X, \beta)^T$ to separate the contributions from components of initial type and the contributions from components of recursion type, we conclude the following Lemmas, whose proofs are word for word the same as the proofs of Lemma 7.5.1 and of the Recursion Lemma 7.5.2 in [9].

Lemma 3.7.1. *Each $(q, \{t_i\})$ -coefficient of S_ν^ε is naturally an element of $K(z)$. Further, this rational function decomposes as a sum of partial simple fractions with denominators either powers of z , or powers of $(z - \alpha)$ with $-\alpha$ a weight of the T -representation on the tangent space $T_\nu \bar{I}_\mu X$ for some $n \in \mathbb{Q}_{>0}$.*

Lemma 3.7.2. *S_ν^ε satisfies the recursion relation*
(3.7.2)

$$S_\nu^\varepsilon(z) = R_\nu^\varepsilon(z) + \sum_{M' \in U(\nu)} q^{\beta_{M'}} \left\langle \frac{\delta_\nu}{z - \alpha_{M'} - \psi_0}, S_{\nu_{M'}, \alpha_{M'}}^\varepsilon \Big|_{z=\alpha_{M'} - \psi_\infty} \right\rangle_{M'}$$

where:

- $\delta_\nu := (i_\nu)_*(\mathbb{1}_\nu)$
- $U(\nu)$ is the set of all unbroken components $M' \subset \mathcal{K}_{0,2}(X, \beta_{M'})^T$ for varying $\beta_{M'}$ for which the first marking lands on ν via the evaluation map.
- $\nu_{M'} \in (\bar{I}_\mu X)^T$ denotes $f(x_2)$ for $((C, x_1, x_2), f) \in M'$.
- ψ_0, ψ_∞ are the nonequivariant Psi classes on M' associated to the markings x_1, x_2 , respectively.
- $R_\nu^\varepsilon(z)$ is the contribution from all components of initial type and has the property that each of its $(q, \{t_i\})$ -coefficients is an element in $K[1/z]$.
- For $\nu \in (\bar{I}_\mu X)^T$, $S_{\nu, \alpha}^\varepsilon(z)$ denotes the part of $S_\nu^\varepsilon(z)$ remaining after the partial fraction terms with poles at $z = \alpha$ are removed.
- The subscript M' for the bracket means the T -virtual localization contribution of the component M' to the virtual intersection number on $\mathcal{K}_{0,2}(X, \beta_{M'})$.

3.7.3. *Uniqueness.* Lemmas parallel to Polynomiality Lemma 7.6.1, Uniqueness Lemma 7.7.1 of [9] hold. We mention the needed minor modifications.

- (1) Instead of X^T , we need to use the set of \mathbb{C} -points of $\bar{I}_\mu X^T$, i.e.,

$$\{(x, g) : x \in X^T(\mathbb{C}), g \in \text{Aut}(x)\}.$$

- (2) In the polynomiality formula in Lemma 7.6.1 of [9], the product between two S_ν^ε now becomes

$$S_\nu^\varepsilon(q, t, z) \iota^*(S_{\nu^{-1}}^\varepsilon(qe^{-zyL_\theta}, t, -z)),$$

where ν^{-1} denotes the point obtained from ν after inverting the band structure, i.e., $\nu^{-1} := (x, g^{-1})$ for $\nu = (x, g)$. Note that $\iota^*S_{\nu^{-1}}^\varepsilon(q, t, z) = i_\nu^*S^\varepsilon(q, t, z) = S_\nu^\varepsilon(q, t, z)$.

- (3) $S_\nu^\varepsilon(q, t, z)$ modulo q does not depend on ε . (This is observed in Remark 3.2.2(3) and replaces condition (5) in Uniqueness Lemma 7.7.1 of [9].)

With these changes in mind, the argument of [9, §7.8, 7.9] applies to the orbifold setting and provides the proof of Theorem 3.6.2.

Remark 3.7.3. (Twisted theories) Let \mathcal{E} be a vector bundle on \mathfrak{X} or, equivalently, a G -equivariant vector bundle on W (for example, one could take $\mathcal{E} = W \times E$, with E a G -representation). Given an invertible \mathbb{G}_m -equivariant characteristic class c , we may define (\mathcal{E}, c) -twisted orbifold ε -quasimap invariants exactly as in [12, §6.2] (here $\mathbb{G}_m = \mathbb{C}^*$ acts by scaling in the fibers of vector bundles). Let $c = Euler$ be the \mathbb{G}_m -equivariant Euler class. We then have the twisted versions of the J^ε -functions and of the S^ε -operators, and as in [9] the proof of Theorem 3.6.2 applies in this context as well.

Let $s \in \Gamma(W, \mathcal{E})^G$ be a regular section and let $Z \in W$ be its G -invariant zero locus. Assume that $Z^{ss} = W^{ss} \cap Z$ is nonsingular. The bundle \mathcal{E} descends to a vector bundle $\underline{\mathcal{E}}$ on X and s descends to a regular section \underline{s} of $\underline{\mathcal{E}}$. The pair $(Y := [Z^{ss}/G], \mathfrak{Y} := [Z/G])$ (or the triple (Z, G, θ)) satisfies the conditions of Theorem 2.3.7 and therefore has its own quasimap theory. (Note that Y is the zero locus of \underline{s} .) If \mathcal{E} is convex (by definition, this means that $H^1(C, [u]^*\mathcal{E}) = 0$ for all genus zero θ -quasimaps $[u] : C \rightarrow \mathfrak{X}$), then in genus zero the $(\mathcal{E}, Euler)$ -twisted quasimap theory of X coincides with the quasimap theory of Y with insertions restricted to Chen-Ruan cohomology classes pulled-back from $\bar{I}_\mu X$. Hence in this situation Theorem 3.6.2 gives the wall-crossing formulas for Y under the same assumptions on the T -action on W .

4. I-FUNCTIONS AND STACKY LOOP SPACES

In this section we introduce (after [11]) a generalization of Givental's small I -function and prove that it lies on the Lagrangian cone of the Gromov-Witten theory of X . We then show that by using certain "stacky loop spaces" of θ -quasimaps to \mathfrak{X} , these new \mathbb{I} -functions can be explicitly computed.

4.1. \mathbb{I} -functions. Denote by $I(t, q, z)$ the J -function for $(0+)$ -quasimap theory $J^{0+}(t, q, z)$ and let

$$I(0, q, z) := \sum q^\beta I_\beta(0, q, z)$$

be its specialization at $t = 0$.

Let $t = \sum_i t_i \gamma_i \in H_T^*(X)$ (in the untwisted sector), with the sum taken only over those γ_i which can be written as a polynomial

$$(4.1.1) \quad p_i(c_1(L_{\eta_{ij}})) = p_i(c_1(L_{\eta_{i1}}, \dots, c_1(L_{\eta_{im_i}}))$$

in divisor classes of the form $c_1(L_{\eta_{ij}})$ for some $\eta_{ij} \in \chi(G)$.

Definition 4.1.1.

$$\mathbb{I}(t, q, z) = \sum_\beta q^\beta \exp\left(\frac{1}{z} \sum_i t_i p_i(c_1(L_{\eta_{ij}}) + \beta(L_{\eta_{ij}})z)\right) I_\beta(0, q, z).$$

The specialization of $\mathbb{I}(t, q, z)$ to

$$t = t_0 \mathbb{1}_X + \sum t_i c_1(L_{\eta_i}) \in H_T^{\leq 2}(X)$$

for $\eta_i \in \chi(G)$ is called *Givental's small I-function* and denoted by $I^{\text{Giv}}(t, q, z)$.

The following is the second main result of the paper.

Theorem 4.1.2. *Assume that the T -action on \underline{X} has isolated T -fixed points and isolated 1-dimensional T -orbits. Then $\mathbb{I}(t, q, z)$ is on the Lagrangian cone of the Gromov-Witten theory of X .*

Proof. The argument is similar to the proof (under the same assumptions on the T -action) of the second part of Theorem 3.6.2.

By Lemma 6.4.1 of [9], there is a unique transformation

$$t \mapsto \tau^{\infty, \varepsilon}(t) \in H_T^*(\bar{I}_\mu X) \otimes_{\mathbb{Q}} K[[\text{Eff}(W, G, \theta)]][[\{t_i\}_i]]$$

and a unique element

$$P^{\infty, \varepsilon}(t, q, z) \in H_T^*(\bar{I}_\mu X) \otimes_{\mathbb{Q}} K[z][[\text{Eff}(W, G, \theta)]][[\{t_i\}_i]]$$

such that

$$\mathbb{I}(t, q, z) = S_{\tau^{\infty, \varepsilon}(t)}^{\infty}(P^{\infty, \varepsilon}(t, q, z)) \text{ modulo } 1/z^2.$$

Now if we let

$$\begin{aligned} S_{1, \nu} &= i_\nu^* \mathbb{I}(t, q, z) \\ S_{2, \nu} &= i_\nu^* S_{\tau^{\infty, \varepsilon}(t)}^{\infty}(P^{\infty, \varepsilon}(t, q, z)). \end{aligned}$$

then it is straightforward to check that the systems $\{S_{i, \nu} \mid \nu \in \bar{I}_\mu X^T\}$, $i = 1, 2$ satisfy all the properties in Uniqueness Lemma 7.7.1 of [9]

(replacing $W//G^T$ by $\bar{I}_\mu X^T$ and condition (5) of Uniqueness Lemma 7.7.1 [9] by $S_{1,\nu} = S_{2,\nu}$ modulo q) so that $S_{1,\nu} = S_{2,\nu}$ for all ν . More precisely, we need to check the conditions (1) – (3) of the Uniqueness Lemma. For $\{S_{2,\nu}\}_\nu$ this is already done in §3.7. Similarly, §3.7 shows that $\{i_\nu^* I(0, q, z)\}_\nu$ satisfies conditions (1) – (3) (since it is the specialization of $\{i_\nu^* S_t^{0+}(P^{0+}(t, q, z))\}_\nu$ at $t = 0$). From this fact and the explicit form of the exponential correcting factor, by using the simple observation

$$(c_1(L_{\eta_{ij}}) + \beta(L_{\eta_{ij}})z)|_{\nu, z=\alpha_{M'}} = (c_1(L_{\eta_{ij}}) + (\beta - \beta_{M'})(L_{\eta_{ij}})z)|_{\nu', z=\alpha_{M'}},$$

it follows by a direct check that conditions (1) – (3) continue to hold for $\{i_\nu^* \mathbb{I}(t, q, z)\}_\nu$. \square

Remark 4.1.3. Alternatively, and better, one may use the geometric arguments of §3.7 to show that $\{i_\nu^* \mathbb{I}(t, q, z)\}_\nu$ satisfies conditions (1) – (3) in the Uniqueness Lemma 7.7.1 of [9]. We explain this briefly. Consider the quasimap theory with weighted markings from [11], extended to orbifold targets using §2.5.5. Fix the stability $(0+, 0+)$, that is, the asymptotic stability with respect to θ and infinitesimally small weights on the weighted markings. For this theory, we have the J -function $\mathbb{J}^{0+, 0+}(\mathbf{t}, q, z)$, the S -operator $\mathbb{S}_{\mathbf{t}}^{0+, 0+}$, and the P -series $P^{0+, 0+}(\mathbf{t}, q, z)$, satisfying the Birkhoff factorization

$$(4.1.2) \quad \mathbb{J}^{0+, 0+}(\mathbf{t}, q, z) = \mathbb{S}_{\mathbf{t}}^{0+, 0+}(P^{0+, 0+}(\mathbf{t}, q, z)).$$

Here $\mathbf{t} \in H_T^*([W/G])$ is a general element. By §5 of [11], the new I -function \mathbb{I} of Definition 4.1.1 is identical with $\mathbb{J}^{0+, 0+}(\mathbf{t}, q, z)$ after the specialization $\mathbf{t} = \sum t_i \tilde{\gamma}_i$ where $\tilde{\gamma}_i$ is the natural lift of $\gamma_i = p_i(c_1(L_{\eta_{ij}}))$ by taking the Chern class of the corresponding line bundles on $[W/G]$. Through this identification, the fact that $\{i_\nu^* \mathbb{I}(t, q, z)\}_\nu$ satisfies conditions (1) – (3) follows from the geometric argument of §3.7 applied for RHS of (4.1.2).

4.2. Stacky loop spaces. In view of (the semi-positive case of) Theorem 3.6.2 and Theorem 4.1.2, it is important to compute explicitly the function $I(0, q, z) = J^{0+}|_{t=0}$. For this purpose, we construct another quasimap graph space. When X is a weighted projective space, this construction already appeared in [14].

Denote by $\mathbb{P}_{a,1}$ the quotient stack $[\mathbb{C}^2 \setminus \{0\}/\mathbb{C}^*]$ where \mathbb{C}^* acts on \mathbb{C}^2 by weights a and 1. Note that $0 := [0, 1]$ is a schematic point, while $\infty := [1, 0] \cong B\mu_a$ is a stacky point for $a > 1$.

For a positive integer a and $\beta \in \text{Hom}_{\mathbb{Z}}(\text{Pic}(\mathfrak{X}), \mathbb{Q})$, define

$$Q_{\mathbb{P}_{a,1}}(X, \beta) \subset \text{Hom}_{\beta}^{\text{rep}}(\mathbb{P}_{a,1}, \mathfrak{X})$$

to be the moduli stack of all θ -quasimaps $[u]$ from $\mathbb{P}_{a,1}$ to $\mathfrak{X} = [W/G]$. This means that $[u] \in Q_{\mathbb{P}_{a,1}}(X, \beta)$ is a representable morphism to \mathfrak{X} , mapping the generic point of $\mathbb{P}_{a,1}$ into X . However, the stacky point $\infty = [1, 0]$ is allowed to be mapped to the unstable locus $\mathfrak{X} \setminus X$.

Proposition 4.2.1. *The stack $Q_{\mathbb{P}_{a,1}}(X, \beta)$ is a DM stack proper over X_0 , equipped with a canonical perfect obstruction theory $R^\bullet \pi_* [u]^* \mathbb{T}_{\mathfrak{X}}^\vee$.*

Proof. Since it is a substack of $\text{Hom}_{\text{Spec } \mathbb{C}}(\mathbb{P}_{a,1}, \mathfrak{X})$, by Proposition 2.11 of [22], it is an Artin stack of locally finite type over \mathbb{C} . By the boundedness Theorem 3.2.4 of [12], it is of finite type over \mathbb{C} . The deformation/obstruction theory is clearly given by $R^\bullet \pi_* [u]^* \mathbb{T}_{\mathfrak{X}}^\vee$ and by the quasimap condition, there is no infinitesimal automorphism so that the stack is a DM stack. The only remaining part is to show the stack is proper. As before, we use the valuative criterion for properness. For $a = 1$, this is known by [12]. For $a > 1$, the argument in [12] applied on étale charts of $\mathbb{P}_{a,1}$ works. \square

The following lemma gives a condition on a which is necessary for the non-emptiness of $Q_{\mathbb{P}_{a,1}}(X, \beta)$.

Lemma 4.2.2. *Let $T(G)$ denote a maximal torus of G . Every morphism $[u] \in \text{Hom}_\beta(\mathbb{P}_{a,1}, \mathfrak{X})$ induces a canonical homomorphism $\tilde{\beta} : \chi(T(G)) \rightarrow \mathbb{Q}$, well-defined up to the Weyl group action on the character group $\chi(T(G))$. Furthermore, $[u]$ is representable if and only if a is the minimal positive integer making $a\tilde{\beta}(\eta) \in \mathbb{Z}$ for all $\eta \in \chi(T(G))$.*

Proof. Every $[u] : \mathbb{P}_{a,1} = [\mathbb{C}^2 \setminus \{0\}/\mathbb{C}^*] \rightarrow \mathfrak{X} = [W/G]$ is the restriction of a unique morphism $[\mathbb{C}^2/\mathbb{C}^*] \rightarrow [W/G]$. The latter induces a morphism $B\mathbb{C}^* \rightarrow BG$ or, equivalently, a group homomorphism $\lambda : \mathbb{C}^* \rightarrow G$ unique up to conjugacy classes. We may assume that λ is factored through the inclusion $T(G) \subset G$. Define $\tilde{\beta} \in \text{Hom}(\chi(T(G)), \mathbb{Q})$ by

$$\tilde{\beta}(\eta) = \frac{\text{the exponent of } (\eta \circ \lambda)}{a} \in \mathbb{Q}.$$

Note that the map $[u]$ is representable if and only if $\lambda|_{\mu_a}$ is a monomorphism, where we identify

$$\mu_a = \langle \xi \rangle \subset \mathbb{C}^*, \quad \xi := e^{2\pi\sqrt{-1}/a}.$$

Let k be the smallest integer for which $\xi^k \in \text{Ker } \lambda|_{\mu_a}$ and $0 < k \leq a$. This means that k is the smallest integer among $0 < k \leq a$ such that $k\tilde{\beta}(\eta) \in \mathbb{Z}$, $\forall \eta$. Thus, $\lambda|_{\mu_a}$ is a monomorphism if and only if a is the minimal positive integer making $a\tilde{\beta}(\eta) \in \mathbb{Z}$, $\forall \eta$. \square

Remark 4.2.3. Let the quasimap $[u] : \mathbb{P}_{a,1} \rightarrow [W/G]$ be given by the data $(\mathbb{P}_{a,1}, P, u)$. By the extension of Grothendieck's theorem (see Theorem 2.4 and Theorem 2.7 in [24]), the principal G -bundle P has a reduction $P_{T(G)}$ to the maximal torus $T(G)$, whose isomorphism class is unique up to the action of the Weyl group. The data $(\mathbb{P}_{a,1}, P_{T(G)}, u)$ gives a quasimap

$$[\widetilde{u}] : \mathbb{P}_{a,1} \longrightarrow [W/T(G)]$$

which lifts $[u]$. It follows that, up to the Weyl group action, the associated numerical class

$$\beta_{[\widetilde{u}]} : \text{Pic}([W/T(G)]) \rightarrow \mathbb{Q}$$

is uniquely determined by $[u]$. It is immediate to see that the homomorphism $\tilde{\beta}$ in Lemma 4.2.2 is the restriction of $\beta_{[\widetilde{u}]}$ to $\chi(T(G))$.

For $\beta \in \text{Eff}(W, G, \theta)$, denote

$$Q_{\mathbb{P}_{\bullet,1}}(X, \beta) = \prod_{1 \leq a \leq e} Q_{\mathbb{P}_{a,1}}(X, \beta)$$

with the induced absolute perfect obstruction theory $(R^\bullet \pi_* [u]^* T_{\mathfrak{X}})^\vee$. Let F_β be the distinguished \mathbb{C}^* -fixed closed substack of $Q_{\mathbb{P}_{\bullet,1}}(X, \beta)$ consisting of elements which have a single base-point of length $\beta(L_\theta)$ at the point $[0, 1]$ of $\mathbb{P}_{a,1}$ (i.e., the class β is exactly supported at $[0, 1]$).

For simplicity we write $QG := QG_{0,\star,\beta}^{0+}(X)$, $Q_{\mathbb{P}_{\bullet,1}} := Q_{\mathbb{P}_{\bullet,1}}(X, \beta)$.

Lemma 4.2.4. (1) *There is a natural isomorphism between an open neighborhood of $F_{\star,0}^{\theta,\beta}$ in the closed substack $(\pi_{\mathbb{P}^1} \circ ev_\star)^{-1}(\infty)$ of QG and an open neighborhood of F_β in $Q_{\mathbb{P}_{\bullet,1}}$, under which $F_{\star,0}^{\theta,\beta} \cong F_\beta$. The isomorphism preserves the $\mathbb{C}^* \times T$ -equivariant perfect obstruction theories.*

(2) *Under the natural isomorphism between $F_{\star,0}^{\theta,\beta}$ and F_β ,*

$$\frac{ev_\star^*(\eta_\infty)[F_{\star,0}^{\theta,\beta}]^{\text{vir}}}{e^{\mathbb{C}^* \times T}(N_{F_{\star,0}^{\theta,\beta}/QG}^{\text{vir}})} = \frac{[F_\beta]^{\text{vir}}}{e^{\mathbb{C}^* \times T}(N_{F_\beta/Q_{\mathbb{P}_{\bullet,1}}}^{\text{vir}})}.$$

Here again the localization residues are taken as sums over the connected components of $F_{\star,0}^{\theta,\beta}$ and of F_β .

Proof. (1): Take the open neighborhood in QG by imposing the condition that the domain curves are irreducible and the open neighborhood in $Q_{\mathbb{P}_{\bullet,1}}$ by requiring that the base points are away from the stacky point $[1, 0] \subset \mathbb{P}_{a,1}$.

(2): We compare the \mathbb{C}^* moving and fixed parts of both obstruction theories. First for QG , we need to look at the fixed part of

$R^\bullet \pi_*([u]_1^* T_{\mathfrak{X}} \boxtimes [u]_2^* T\mathbb{P}^1)$ and the fixed part of the infinitesimal automorphism/deformation of (C, x_*) . Altogether its contribution coincides with the fixed part of $R^\bullet [u]^* T_{\mathfrak{X}}$. The Euler class of the moving part of them altogether becomes the Euler class of the moving part of $R^\bullet [u]^* T_{\mathfrak{X}}$ divided by $(-z)$. \square

By Lemma 3.3.4 and Lemma 4.2.4, we obtain the following.

Proposition 4.2.5.

$$I(0, q, z) = \mathbb{1}_X + \sum_{\beta \neq 0} q^\beta (\tilde{e}v_\star)_* \frac{[F_\beta]^{\text{vir}}}{e^{\mathbb{C}^* \times T} (N_{F_\beta/Q_{\mathbb{P}^{\bullet}, 1}}^{\text{vir}})}.$$

5. TORIC ORBIFOLDS

In this section we make Theorem 4.1.2 completely explicit for toric orbifolds by calculating the localization residues $I_\beta(0, q, z)$ via stacky loop spaces. For toric manifolds this is a well known calculation with Euler sequences, due to Givental [16]. For the convenience of the reader we present its extension to the orbifold case. As a result of this calculation, the Mirror Theorem for toric orbifolds, recently proved by different methods in [13], becomes a special case of Theorem 4.1.2.

5.1. Set-up. Let G be the algebraic torus $(\mathbb{C}^*)^r$, $r \geq 0$. Fix a finite collection $[N]$ of (not necessarily distinct) characters of G . In this section, we consider the case

$$W = \bigoplus_{\rho \in [N]} \mathbb{C}_\rho, \quad G = (\mathbb{C}^*)^r.$$

Fix a character θ of G and assume that $W^{ss} := W^{ss}(\theta) = W^s(\theta)$ as before. For a character ρ , the associated line bundle on $\mathfrak{X} := [W/G]$ will be denoted by L_ρ . Let π_i be the i -th ‘‘standard’’ character

$$\pi_i : G = (\mathbb{C}^*)^r \longrightarrow \mathbb{C}^*,$$

coming from the i -th projection. As the character group of G is the free abelian group generated by π_i , there are unique integers $a_{i,\rho}$ making $\rho = \sum_i a_{i,\rho} \pi_i$.

The GIT stack quotient $X := [W^{ss}/G]$ is a toric DM stack (in the sense of [5]), with quasi-projective coarse moduli space. Since it is a global quotient stack by an abelian group, it is known that

$$I_\mu X = \prod_{g \in G} [(W^{ss})^g/G], \quad \text{and hence} \quad \bar{I}_\mu X = \prod_{g \in G} [(W^{ss})^g/(G/\langle g \rangle)].$$

Let $T = (\mathbb{C}^*)^{[N]}$ be the big torus with the standard action on W . There are only finitely many fixed points and finitely many 1-dimensional

orbits on \underline{X} under the induced T action. Therefore Theorems 3.6.2 and 4.1.2 apply to the triple (W, G, θ) .

In what follows, we will denote D_ρ the hyperplane of W associated to ρ as well as the corresponding T -equivariant divisor classes of $[W/H]$ (or even its restriction to various substacks of $[W/H]$), whenever H is an algebraic group acting on W with the hyperplane D_ρ being invariant.

5.2. Explicit description of $\mathrm{Hom}_\beta(\mathbb{P}_{a,1}, [W/G])$. Let $\mathcal{S}ch_{\mathbb{C}}$ be the category of schemes over $\mathrm{Spec}\mathbb{C}$ and let $U := \mathbb{C}^2 \setminus \{0\}$.

Since $\mathrm{Pic}U \cong \mathrm{Pic}\mathbb{C}^2$ is trivial, any line bundle on $\mathbb{P}_{a,1}$ is obtained from the Borel's mixed construction $\mathcal{O}_{\mathbb{P}_{a,1}}(m) := [U \times \mathbb{C}_m / \mathbb{C}^*]$ for some weight $m \in \mathbb{Z}$ of \mathbb{C}^* . The degree of $\mathcal{O}_{\mathbb{P}_{a,1}}(m)$ is $m/a \in \mathbb{Q}$.

We find an explicit description of Hom-stack $\mathrm{Hom}_\beta(\mathbb{P}_{a,1}, \mathfrak{X})$. The groupoid fiber over $S \in \mathcal{S}ch_{\mathbb{C}}$ of the category $\mathrm{Hom}_\beta(\mathbb{P}_{a,1}, \mathfrak{X})$ is equivalent to the category described as follows:

- Objects are collections

$$(\mathcal{L}_i, i \in [r]; u_\rho, \rho \in [N])$$

consisting of \mathbb{C}^* -equivariant line bundles \mathcal{L}_i on $S \times U$, $i = 1, \dots, r$, together with \mathbb{C}^* -invariant sections u_ρ of

$$\mathcal{L}_\rho := \otimes_i \mathcal{L}_i^{\otimes a_i \rho}.$$

By triviality of $\mathrm{Pic}U$, the \mathbb{C}^* -equivariant line bundle \mathcal{L}_i is determined uniquely by an integer weight of \mathbb{C}^* and a line bundle M_i on S . For the collection to give a map of class β , the weight must be $a\beta(L_{\pi_i}) \in \mathbb{Z}$:

$$\mathcal{L}_i = M_i \boxtimes (U \times \mathbb{C}_{a\beta(L_{\pi_i})}).$$

- Arrows from $(\mathcal{L}_i, i \in [r]; u_\rho, \rho \in [N])$ to $(\mathcal{L}'_i, i \in [r]; u'_\rho, \rho \in [N])$ are collections of isomorphisms $(\varphi_i \in \Gamma(S \times U, \mathcal{L}_i^\vee \otimes \mathcal{L}'_i)^{\mathbb{C}^*}, i \in [r])$ for which u_ρ corresponds to u'_ρ for every ρ .

Consider the graded ring $\mathbb{C}[x, y]$ with $\deg x = a$, $\deg y = 1$ and denote by $\mathbb{C}[x, y]_m$ its \mathbb{C} -subspace of degree m . Pushing-forward along U , together with the fact that $H^0(U, \mathcal{O}_U) = H^0(\mathbb{C}^2, \mathcal{O}_{\mathbb{C}^2}) = \mathbb{C}[x, y]$, we may regard u_ρ canonically as an element of

$$\Gamma(S, M_\rho) \otimes_{\mathbb{C}} \mathbb{C}[x, y]_{a\beta(L_\rho)},$$

where $M_\rho := \otimes_i M_i^{\otimes a_i \rho}$.

Hence the groupoid fiber above is equivalent to the category whose objects are $(M_i, u_\rho \in \Gamma(S, M_\rho) \otimes_{\mathbb{C}} \mathbb{C}[x, y]_{a\beta(L_\rho)}, \rho \in [N])$ and whose arrows from $(M_i, u_\rho \in \Gamma(S, M_\rho) \otimes_{\mathbb{C}} \mathbb{C}[x, y]_{a\beta(L_\rho)}, \rho \in [N])$ to $(M'_i, u'_\rho \in$

$\Gamma(S, M'_i) \otimes \mathbb{C}[x, y]_{a\beta(L_\rho)}$, $\rho \in [N]$) are collections of \mathcal{O}_S -module isomorphisms $M_i \rightarrow M'_i$, $i \in [r]$, compatible with u_ρ, u'_ρ .

Given a and β , consider the finite dimensional vector space

$$W_\beta^a := \bigoplus_{\rho \in [N]} \mathbb{C}[x, y]_{a\beta(L_\rho)}$$

with the G action given by the direct sum of the diagonal G action on $\mathbb{C}[x, y]_{a\beta(L_\rho)}$ by the weight ρ , so that $\mathbb{C}[x, y]_{a\beta(L_\rho)} \cong \bigoplus \mathbb{C}_\rho$. In particular, when $\beta = 0$, we recover $W_0 = W$ with the original G action. Now the conclusion of the equivalent descriptions of the groupoids above can be stated as follows.

Proposition 5.2.1. *The following stacks are all equivalent:*

$$\mathrm{Hom}_\beta(\mathbb{P}_{a,1}, \mathfrak{X}) \cong \mathrm{Hom}_\beta([\mathbb{C}^2/\mathbb{C}^*], \mathfrak{X}) \cong [W_\beta^a/G].$$

Convention: From now on we let a be the minimal positive integer associated to β by Lemma 4.2.2 and will write W_β for W_β^a .

Corollary 5.2.2. *The stack $Q_{\mathbb{P}_{a,1}}(X, \beta)$ is equivalent to the smooth quotient stack $[W_\beta^{ss}(\theta)/G]$.*

Proof. By Proposition 5.2.1 and Lemma 4.2.2, $[u] \in \mathrm{Hom}_\beta(\mathbb{P}_{a,1}, \mathfrak{X})$ is a \mathbb{C} -point of $Q_{\mathbb{P}_{a,1}}(X, \beta)$ if and only if $u_\rho(\zeta_0, \zeta_1) \in W^{ss}(\theta)$ for general points $(\zeta_0, \zeta_1) \in \mathbb{C}^2 \setminus \{(0, 0)\}$. It is easy to see that the latter condition is equivalent to $(u_\rho)_\rho \in W_\beta^{ss}(\theta)$. \square

Recall the perfect obstruction theory $R^\bullet \pi_*[u]^* \mathbb{T}_{[W/G]}$ for $Q_{\mathbb{P}_{a,1}}(X, \beta)$ as in §4.2. We can describe the complex more explicitly since $[W_\beta^{ss}/G]$ is smooth. Consider the generalized Euler sequence (see [12, §5.1]), i.e., the distinguished triangle

$$\mathfrak{g} \times_G W \rightarrow W \times_G W \rightarrow \mathbb{T}_{[W/G]}.$$

on $[W/G]$. It induces an exact sequence on $[W_\beta^{ss}/G]$

$$0 \rightarrow \mathcal{O}^{\oplus[r]} \rightarrow \bigoplus_{\rho \in [N]} \pi_*[u]^* L_\rho \rightarrow \mathbb{T}_{[W_\beta^{ss}/G]} \rightarrow 0,$$

and the obstruction vector bundle $\bigoplus_{\rho \in [N]} R^1 \pi_*[u]^* L_\rho$ defining the virtual fundamental class of $[W_\beta^{ss}/G] \cong Q_{\mathbb{P}_{a,1}}(X, \beta)$.

5.3. The virtual normal bundle. Recall the \mathbb{C}^* action on the coarse moduli \mathbb{P}^1 , given as $t \cdot [\zeta_0, \zeta_1] = [t\zeta_0, \zeta_1]$. This induces an action on $[W_\beta^{ss}/G]$. Define

$$(5.3.1) \quad Z_\beta := \bigoplus_{\rho \in [N], \beta(L_\rho) \in \mathbb{Z}_{\geq 0}} \mathbb{C} \cdot x^{\beta(L_\rho)} \subset W_\beta.$$

Under the natural identification of Z_β with the sub- G -representation $\bigoplus_{\rho \in [N], \beta(L_\rho) \in \mathbb{Z}_{\geq 0}} \mathbb{C}_\rho$ of W , we have

$$Z_\beta = \bigcap_{\rho: \beta(L_\rho) < 0 \text{ or } \beta(L_\rho) \notin \mathbb{N}} D_\rho.$$

The \mathbb{C}^* -fixed component $[Z_\beta/G]$ of $\text{Hom}_\beta(\mathbb{P}_{a,1}, X)$ is distinguished in the sense that

$$F_\beta \cong [Z_\beta/G] \cap [W_\beta^{ss}/G]$$

under the identification from Corollary 5.2.2. Let $Z_\beta^{ss} := Z_\beta \cap W_\beta^{ss}$.

Proposition 5.3.1.

$$\begin{aligned} e^{\mathbb{C}^* \times T}(N_{F_\beta/QG_{\mathbb{P}_{a,1}}}^{\text{vir}}) &= e^{\mathbb{C}^* \times T}(N_{[Z_\beta^{ss}/G]/[W_\beta^{ss}/G]}) \\ &= \frac{\prod_{\rho: \beta(L_\rho) > 0} \prod_{0 \leq \nu < \beta(L_\rho)} (D_\rho + (\beta(L_\rho) - \nu)z)}{\prod_{\rho: \beta(L_\rho) < 0} \prod_{\lfloor \beta(L_\rho) + 1 \rfloor \leq \nu < 0} (D_\rho + (\beta(L_\rho) - \nu)z)} \end{aligned}$$

and $[F_\beta]^{\text{vir}} = [F_\beta] = [Z_\beta^{ss}/G]$. (In the above formula the index ν in the products runs over integers and $\lfloor \cdot \rfloor$ denotes the round-down of a rational number.)

Proof. First note that $\pi_*[u]^*L_\rho = \mathcal{O}(D_\rho) \otimes H^0(\mathbb{P}_{a,1}, \mathcal{O}(a\beta(L_\rho)))$. The \mathbb{C}^* -representation space $H^0(\mathbb{P}_{a,1}, \mathcal{O}(a\beta(L_\rho)))$ (for $\beta(L_\rho) \geq 0$) has the basis

$$y^{a\beta(L_\rho)}, y^{a(\beta(L_\rho)-1)}x, \dots, y^{a(\beta(L_\rho)-\lfloor \beta(L_\rho) \rfloor)}x^{\lfloor \beta(L_\rho) \rfloor},$$

therefore its \mathbb{C}^* weights are $\beta(L_\rho), \beta(L_\rho) - 1, \dots, \beta(L_\rho) - \lfloor \beta(L_\rho) \rfloor$. This explains the numerator of $e^{\mathbb{C}^* \times T}(N_{F_\beta/QG_{\mathbb{P}_{a,1}}}^{\text{vir}})$.

The obstruction bundle $\bigoplus_\rho R^1\pi_*[u]^*L_\rho$ becomes

$$\bigoplus_{\rho: \beta(L_\rho) \in \mathbb{Q}_{<0}} \mathcal{O}(D_\rho) \otimes H^1(\mathbb{P}_{a,1}, \mathcal{O}(a\beta(L_\rho))).$$

A \mathbb{C}^* -eigenbasis of the cohomology space $H^1(\mathbb{P}_{a,1}, \mathcal{O}(a\beta(L_\rho)))$ can be computed by taking Čech 1-cocycles with respect to the étale covering of $\mathbb{P}_{a,1}$:

$$y^{a(\beta(L_\rho)+1)}x^{-1}, y^{a(\beta(L_\rho)+2)}x^{-2}, \dots, y^{a(\beta(L_\rho)-\lfloor \beta(L_\rho)+1 \rfloor)}x^{\lfloor \beta(L_\rho)+1 \rfloor}.$$

Therefore the weights are $\beta(L_\rho) - \nu$ with $\lfloor \beta(L_\rho) + 1 \rfloor \leq \nu < 0$, $\nu \in \mathbb{Z}$. This is the denominator of $e^{\mathbb{C}^* \times T}(N_{F_\beta/QG_{\mathbb{P}_{a,1}}}^{\text{vir}})$.

Finally, the obstruction bundle defining the virtual fundamental class of the smooth stack $[W_\beta^{ss}(\theta)/G] = [W_\beta^s(\theta)/G]$, when restricted to F_β has no \mathbb{C}^* -fixed part. We conclude that $[F_\beta]^{\text{vir}} = [F_\beta]$. \square

Let $g_\beta = (e^{2\pi\sqrt{-1}\beta(L_\rho)})_\rho \in G$. By the proof of Lemma 4.2.2, note that the connected component into which F_β lands under the evaluation map at \star and its associated component in the unrigidified cyclotomic inertia stack are respectively

$$(5.3.2) \quad \begin{aligned} & [(W^{ss} \cap \bigcap_{\rho:\beta(\rho) \notin \mathbb{Z}} D_\rho)/(G/\langle g_\beta \rangle)] \text{ and} \\ & [(W^{ss})^{g_\beta}/G] = [(W^{ss} \cap \bigcap_{\rho:\beta(\rho) \notin \mathbb{Z}} D_\rho)/G]. \end{aligned}$$

The evaluation map ev_\star is factored through $[(W^{ss})^{g_\beta}/G]$ and makes $F_\beta = [Z_\beta^{ss}/G]$ as a closed substack of $[(W^{ss})^{g_\beta}/G]$, whose normal bundle yields, by (5.3.1) and (5.3.2),

$$(5.3.3) \quad e^{\mathbb{C}^* \times T}(N_{[Z_\beta^{ss}/G]/[(W^{ss})^{g_\beta}/G]}) = e^{\mathbb{C}^* \times T}(N_{Z_\beta/W^{g_\beta}})|_{[Z_\beta^{ss}/G]} = \prod_{\rho:\beta(L_\rho) \in \mathbb{Z}_{<0}} D_\rho.$$

Theorem 5.3.2.

$$I(0, q, z) = \sum_{\beta \in \text{EH}(W, G, \theta)} q^\beta \frac{\prod_{\rho:\beta(L_\rho) < 0} \prod_{\beta(L_\rho) \leq \nu < 0} (D_\rho + (\beta(L_\rho) - \nu)z)}{\prod_{\rho:\beta(L_\rho) > 0} \prod_{0 \leq \nu < \beta(L_\rho)} (D_\rho + (\beta(L_\rho) - \nu)z)} \mathbb{1}_{g_\beta^{-1}}$$

where $\mathbb{1}_{g_\beta^{-1}}$ is the fundamental class of $[(W^{ss})^{g_\beta^{-1}}/(G/\langle g_\beta^{-1} \rangle)]$.

Proof. This is immediate from Proposition 4.2.5, Proposition 5.3.1, and equation (5.3.3). The factor \mathbf{r} disappears since the rigidification ϖ has degree \mathbf{r}^{-1} . \square

Corollary 5.3.3. *For a semi-positive triple $(W = \mathbb{C}^{[N]}, G = (\mathbb{C}^*)^r, \theta)$, $I(0, q, z)$ takes form*

$$\mathbb{1}_X + \frac{I_1(q)}{z} + O(1/z^2), \quad I_1(q) \in (q\Lambda_K)H_T^{\leq 2}(X, \Lambda_K).$$

Proof. Let $\beta \neq 0$ so that for some ρ , $\beta(L_\rho) \neq 0$. Note that $\mathbb{1}_{g_\beta^{-1}}$ is the fundamental class $\mathbb{1}_X$ of the untwisted sector if and only if $\beta(L_\rho) \in \mathbb{Z}$ for all ρ . In this case, the argument in the proof of Lemma 5.9.1 in [10] works. If $\mathbb{1}_{g_\beta^{-1}}$ is the fundamental class of a twisted sector, then the power of $1/z$ appearing in q^β becomes

$$(5.3.4) \quad \begin{aligned} & \sum_{\rho \in [N]} [\beta(L_\rho)] + \# \text{ negative integers } \beta(L_\rho) \\ & > \beta(\det \mathbb{T}_{\mathfrak{X}}) + \# \text{ negative integers } \beta(L_\rho) \\ & \geq \beta(\det \mathbb{T}_{\mathfrak{X}}) \geq 0. \end{aligned}$$

This implies that $I(0, q, z) = \mathbb{1}_X + O(q)$. The two inequalities in (5.3.4) cannot be equalities at the same time since $\beta \neq 0$. Therefore there is no twisted sector part in the $1/z$ -coefficient of the small I-function $I(0, q, z)$. \square

Let

$$\{\gamma_0 = \mathbb{1}_X, \gamma_1, \dots, \gamma_s, \gamma_{s+1}, \dots, \gamma_l\}$$

be the part of the basis of Chen-Ruan cohomology corresponding to the untwisted sector $H_T^*(X)$, with $\{\gamma_1, \dots, \gamma_s\}$ basis for $H_T^2(X)$. In the toric case, we may write each γ_i as a polynomial in T -equivariant first Chern classes of line bundles coming from characters of G , as in (4.1.1). In particular, for the divisors $\gamma_i, i = 1, \dots, s$ we write $\gamma_i = c_1(L_{\eta_i})$.

Corollary 5.3.4. (1) *Let $t = \sum_{i=0}^l t_i \gamma_i = \sum_{i=0}^l t_i p_i(c_1(L_{\eta_{ij}}))$ be a general element of $H_T^*(X)$. Then*

$$\mathbb{I}(t, q, z) = \sum_{\beta} q^{\beta} \exp\left(\frac{1}{z} \sum_{i=0}^l t_i p_i(c_1(L_{\eta_{ij}}) + \beta(L_{\eta_{ij}})z)\right) \frac{\prod_{\rho: \beta(L_{\rho}) < 0} \prod_{\beta(L_{\rho}) \leq \nu < 0} (D_{\rho} + (\beta(L_{\rho}) - \nu)z)}{\prod_{\rho: \beta(L_{\rho}) > 0} \prod_{0 \leq \nu < \beta(L_{\rho})} (D_{\rho} + (\beta(L_{\rho}) - \nu)z)} \mathbb{1}_{g_{\beta}^{-1}}$$

is on the Lagrangian cone of the Gromov-Witten theory of X .

(2) *For $t = t_0 \mathbb{1}_X + \sum_{i=1}^s t_i \gamma_i \in H_T^0(X) \oplus H_T^2(X)$,*

$$I^{\text{Giv}}(t, q, z) = \sum_{\beta \in \text{Eff}(W, G, \theta)} q^{\beta} e^{(t_0 \mathbb{1}_X + \sum_{i=1}^s t_i (\gamma_i + \beta(L_{\eta_i})z))/z} \frac{\prod_{\rho: \beta(L_{\rho}) < 0} \prod_{\beta(L_{\rho}) \leq \nu < 0} (D_{\rho} + (\beta(L_{\rho}) - \nu)z)}{\prod_{\rho: \beta(L_{\rho}) > 0} \prod_{0 \leq \nu < \beta(L_{\rho})} (D_{\rho} + (\beta(L_{\rho}) - \nu)z)} \mathbb{1}_{g_{\beta}^{-1}}$$

is on the Lagrangian cone of the Gromov-Witten theory of X .

(3) *If $(W = \mathbb{C}^{[N]}, G = (\mathbb{C}^*)^r, \theta)$ is a semi-positive triple,*

$$J^{\infty}(t + I_1(q), q, z) = I^{\text{Giv}}(t, q, z), \text{ for } t \in H_T^{\leq 2}(X).$$

Furthermore, the so-called mirror map $t \mapsto t + I_1(q)$ is expressed via two-pointed $(0+)$ -quasimap invariants as

$$t + I_1(q) = t + \sum_{i=0}^s \sum_{\beta \neq 0} q^{\beta} \gamma_i \langle \gamma^i, \mathbb{1}_X \rangle_{0,2,\beta}^{0+},$$

Proof. (1) is immediate from Theorem 4.1.2 and Theorem 5.3.2, and (2) is a specialization of (1).

(3) follows from (2), Theorem 3.5.2 (3b), and Corollary 5.3.3. \square

In particular, part (3) of the above Corollary proves Conjecture 4.3 of [18] after the identification of $H_T^*(\bar{I}_\mu X)$ with $H_T^*(I_\mu X)$ by the pullback ϖ^* . (Of course, the interpretation of the mirror map as a generating series of quasimap invariants is new.)

Part (2) of the Corollary is precisely the main result of [13]. Note that the notion of “ S -extended I -function” from [13] corresponds in our terminology to the Givental small I -function for a different GIT presentation of the geometric target X .

REFERENCES

- [1] D. Abramovich, T. Graber, A. Vistoli, *Gromov-Witten theory of Deligne-Mumford stacks*, Amer. J. Math. **130** (2008), no. 5, 1337–1398.
- [2] D. Abramovich, M. Olsson, A. Vistoli, *Twisted stable maps to tame Artin stacks*, J. Algebraic Geom. **20** (2011), 399–477.
- [3] D. Abramovich, A. Vistoli, *Compactifying the space of stable maps*, J. Amer. Math. Soc., **15** (2002), no. 1, 27–75.
- [4] K. Behrend, B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128** (1997), no. 1, 45–88.
- [5] L. Borisov, L. Chen, G. Smith, *The orbifold Chow ring of toric Deligne-Mumford stacks*, J. Amer. Math. Soc. **18** (2005), no 1, 193–215.
- [6] C. Cadman, *Using stacks to impose tangency conditions on curves*, Amer. J. Math. **129** (2007), no. 2, 405–427.
- [7] W. Chen, Y. Ruan, *Orbifold Gromov-Witten theory*, in “Orbifolds in mathematics and physics (Madison, WI, 2001)”, 25–85, Contem. Math. **310**, Amer. Math. Soc., 2002.
- [8] I. Ciocan-Fontanine, B. Kim, *Moduli stacks of stable toric quasimaps*, Adv. Math. **225** (2010), no. 6, 3022–3051.
- [9] I. Ciocan-Fontanine, B. Kim, *Wall-crossing in genus zero quasimap theory and mirror maps*, to appear in Algebraic Geometry, arXiv:1304.7056.
- [10] I. Ciocan-Fontanine, B. Kim, *Higher genus quasimap wall-crossing for semi-positive targets*, arXiv:1308.6377v1.
- [11] I. Ciocan-Fontanine, B. Kim, *Big I-functions*, to appear in Proceedings of the conference on the occasion of Mukai’s 60th birthday, arXiv:1401.7417.
- [12] I. Ciocan-Fontanine, B. Kim, D. Maulik, *Stable quasimaps to GIT quotients*, J. of Geometry and Physics **75** (2014), 17–47.
- [13] T. Coates, A. Corti, H. Iritani, H-H. Tseng, *A mirror theorem for toric stacks*, arXiv:1310.4163.
- [14] T. Coates, A. Corti, Y-P. Lee, H-H. Tseng, *The quantum orbifold cohomology of weighted projective spaces*, Acta Math. **202** (2009), no. 2, 139–193.
- [15] A. Deopurkar, *Compactifications of Hurwitz spaces*, to appear in Internat. Math. Res. Notices, arXiv:1206.4535.
- [16] A. Givental, *A mirror theorem for toric complete intersections*, in “Topological field theory, primitive forms and related topics (Kyoto, 1996)”, Progr. Math., 160, Birkhäuser Boston, Boston, MA, 1998, 141–175.
- [17] T. Graber, R. Pandharipande, *Localization of virtual classes*, Invent. Math. **135** (1999), no. 2, 487–518.

- [18] H. Iritani, *An integral structure in quantum cohomology and mirror symmetry for toric orbifolds*, Adv. Math. **222** (2009), no. 3, 1016–1079.
- [19] S. Keel, S. Mori, *Quotients by groupoids*, Ann. of Math. (2) **145** (1997), no. 1, 193–213.
- [20] A. King, *Moduli of representations of finite-dimensional algebras*, Quart. J. Math. Oxford Ser. (2) **45** (1994), no. 180, 515–530.
- [21] A. Kresch, A. Vistoli, *On coverings of Deligne-Mumford stacks and surjectivity of the Brauer map*, Bull. London Math. Soc. **36** (2004), no. 2, 188–192.
- [22] M. Lieblich, *Remarks on the stack of coherent algebras*, Int. Math. Res. Not. 2006, Art. ID 75273, 12 pp.
- [23] A. Marian, D. Oprea, R. Pandharipande, *The moduli space of stable quotients*, Geometry & Topology **15** (2011), 1651–1706.
- [24] J. Martens, M. Thaddeus, *Variations on a theme of Grothendieck*, arXiv:1210.8161.
- [25] A. Mustață, A. Mustață, *Intermediate moduli spaces of stable maps*, Invent. Math. **167** (2007), no. 1, 47–90.
- [26] M. Olsson, *(Log) twisted curves*, Composito Math. **143** (2007), no. 2, 476–494.
- [27] Y. Toda, *Moduli spaces of stable quotients and wall-crossing phenomena*, Compositio Math. **147** (2011), 1479–1518.
- [28] H-H. Tseng, *Orbifold quantum Riemann-Roch, Lefschetz and Serre*, Geom. Topol. **14** (2010), no. 1, 1–81.

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGIRO, DONGDAEMUN-GU, SEOUL, 130-722, KOREA
E-mail address: daewoongc@kias.re.kr

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA, 206 CHURCH ST. SE, MINNEAPOLIS MN, 55455, AND
 SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGIRO, DONGDAEMUN-GU, SEOUL, 130-722, KOREA
E-mail address: ciocan@math.umn.edu

SCHOOL OF MATHEMATICS, KOREA INSTITUTE FOR ADVANCED STUDY, 85 HOEGIRO, DONGDAEMUN-GU, SEOUL, 130-722, KOREA
E-mail address: bumsig@kias.re.kr