

Name: _____

Signature: _____

Math 5651 (V. Reiner) Final Exam
Tuesday, May 10, 2016

This is a 120 minute exam. No books, notes, calculators, cell phones, watches or other electronic devices are allowed. You can leave answers as fractions, with binomial or multinomial coefficients, and Gamma function values $\Gamma(\alpha)$ unevaluated.

There are a total of 100 points. To get full credit for a problem you must show the details of your work. Answers unsupported by an argument will get little credit. Do all of your calculations on this test paper.

Problem	Score
1.	_____
2.	_____
3.	_____
4.	_____
5.	_____
Total:	_____

Reminders:

$$\Pr(A_1 \cup \dots \cup A_n) = \sum_{k=1}^n (-1)^{k-1} \sum_{1 \leq i_1 < \dots < i_k \leq n} \Pr(A_{i_1} \cap \dots \cap A_{i_k})$$

$$S = \sqcup_{i=1}^n B_i \Rightarrow \Pr(A) = \sum_{i=1}^n \Pr(A \cap B_i) = \sum_{i=1}^n \Pr(A|B_i)\Pr(B_i) \text{ and } \Pr(B_i|A) = \Pr(A|B_i)\Pr(B_i)/\Pr(A)$$

$$\text{cdf } F(x) := \Pr(X \leq x), \text{ while pdf } f(x) = \frac{\partial}{\partial x} F(x), \text{ and } g_1(x|y) = f(x, y)/f_2(y) \text{ with } f_2(y) = \int_{x=-\infty}^{x=+\infty} f(x, y) dx$$

$$\text{When } \underline{Y} = \underline{r}(\underline{X}) \Leftrightarrow \underline{X} = \underline{s}(\underline{Y}), \text{ then } f(\underline{x}), g(\underline{y}) \text{ satisfy } g(\underline{y}) = f(\underline{s}(\underline{y})) \cdot |J| \text{ where } J := \det \left(\frac{\partial s_i}{\partial y_j} \right)$$

$$\mathbf{E}X = \int_{-\infty}^{+\infty} x f(x) dx, \text{ and } \text{var}(X) = E(X - EX)^2 = E(X^2) - (EX)^2, \text{ with } \sigma(X) := +\sqrt{\text{var}(X)}$$

$$\text{cov}(X, Y) = E((X - EX)(Y - EY)) = E(XY) - EX \cdot EY = \sigma_X \sigma_Y \rho(X, Y)$$

$$\text{var} \left(\sum_{i=1}^n X_i \right) = \sum_{i=1}^n \text{var}(X_i) + 2 \sum_{1 \leq i < j \leq n} \text{cov}(X_i, X_j)$$

$$\Gamma(\alpha) := \int_{x=0}^{x=\infty} x^{\alpha-1} e^{-x} dx \text{ for } \alpha > 0, \text{ and } \Gamma(\alpha + 1) = \alpha \Gamma(\alpha) \text{ for } \alpha > 1, \text{ with } \Gamma(1) = 1$$

Discrete Distributions

	Bernoulli with parameter p	Binomial with parameters n and p
p.f.	$f(x) = p^x(1-p)^{1-x}$, for $x = 0, 1$	$f(x) = \binom{n}{x} p^x(1-p)^{n-x}$, for $x = 0, \dots, n$
Mean	p	np
Variance	$p(1-p)$	$np(1-p)$
m.g.f.	$\psi(t) = pe^t + 1 - p$	$\psi(t) = (pe^t + 1 - p)^n$

	Uniform on the integers a, \dots, b	Hypergeometric with parameters A, B , and n
p.f.	$f(x) = \frac{1}{b-a+1}$, for $x = a, \dots, b$	$f(x) = \frac{\binom{A}{x}\binom{B}{n-x}}{\binom{A+B}{n}}$, for $x = \max\{0, n-b\}, \dots, \min\{n, A\}$
Mean	$\frac{b+a}{2}$	$\frac{nA}{A+B}$
Variance	$\frac{(b-a)(b-a+2)}{12}$	$\frac{nAB}{(A+B)^2} \frac{A+B-n}{A+B-1}$
m.g.f.	$\psi(t) = \frac{e^{(b+1)t} - e^{at}}{(e^t - 1)(b-a+1)}$	Nothing simpler than $\psi(t) = \sum_x f(x)e^{tx}$

	Geometric with parameter p	Negative binomial with parameters r and p
p.f.	$f(x) = p(1-p)^x$, for $x = 0, 1, \dots$	$f(x) = \binom{r+x-1}{x} p^r(1-p)^x$, for $x = 0, 1, \dots$
Mean	$\frac{1-p}{p}$	$\frac{r(1-p)}{p}$
Variance	$\frac{1-p}{p^2}$	$\frac{r(1-p)}{p^2}$
m.g.f.	$\psi(t) = \frac{p}{1-(1-p)e^t}$, for $t < \log(1/[1-p])$	$\psi(t) = \left(\frac{p}{1-(1-p)e^t}\right)^r$, for $t < \log(1/[1-p])$

	Poisson with mean λ	Multinomial with parameters n and (p_1, \dots, p_k)
p.f.	$f(x) = e^{-\lambda} \frac{\lambda^x}{x!}$, for $x = 0, 1, \dots$	$f(x_1, \dots, x_k) = \binom{n}{x_1, \dots, x_k} p_1^{x_1} \dots p_k^{x_k}$, for $x_1 + \dots + x_k = n$ and all $x_i \geq 0$
Mean	λ	$E(X_i) = np_i$, for $i = 1, \dots, k$
Variance	λ	$\text{Var}(X_i) = np_i(1-p_i)$, $\text{Cov}(X_i, X_j) = -np_i p_j$, for $i, j = 1, \dots, k$
m.g.f.	$\psi(t) = e^{\lambda(e^t - 1)}$	Multivariate m.g.f. can be defined, but is not defined in this text.

Continuous Distributions

	Beta with parameters α and β	Uniform on the interval $[a, b]$
p.d.f.	$f(x) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1}(1-x)^{\beta-1}$, for $0 < x < 1$	$f(x) = \frac{1}{b-a}$, for $a < x < b$
Mean	$\frac{\alpha}{\alpha+\beta}$	$\frac{a+b}{2}$
Variance	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$	$\frac{(b-a)^2}{12}$
m.g.f.	Not available in simple form	$\psi(t) = \frac{e^{-at} - e^{-bt}}{t(b-a)}$

	Exponential with parameter β	Gamma with parameters α and β
p.d.f.	$f(x) = \beta e^{-\beta x}$, for $x > 0$	$f(x) = \frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x}$, for $x > 0$
Mean	$\frac{1}{\beta}$	$\frac{\alpha}{\beta}$
Variance	$\frac{1}{\beta^2}$	$\frac{\alpha}{\beta^2}$
m.g.f.	$\psi(t) = \frac{\beta}{\beta-t}$, for $t < \beta$	$\psi(t) = \left(\frac{\beta}{\beta-t}\right)^\alpha$, for $t < \beta$

	Normal with mean μ and variance σ^2	Bivariate normal with means μ_1 and μ_2 , variances σ_1^2 and σ_2^2 , and correlation ρ
p.d.f.	$f(x) = \frac{1}{(2\pi)^{1/2}\sigma} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right)$	Formula is too large to print here. See Eq. (5.10.2) on page 338.
Mean	μ	$E(X_i) = \mu_i$, for $i = 1, 2$
Variance	σ^2	Covariance matrix: $\begin{pmatrix} \sigma_1^2 & \rho\sigma_1\sigma_2 \\ \rho\sigma_1\sigma_2 & \sigma_2^2 \end{pmatrix}$
m.g.f.	$\psi(t) = \exp\left(\mu t + \frac{t^2\sigma^2}{2}\right)$	Bivariate m.g.f. can be defined, but is not defined in this text.

Problem 1. (20 points) Assume that each of the functions $f(k)$ or $f(x)$ below defines a p.f. or p.d.f., and find the unknown constant c .

a. (4 points) $f(k) = c \frac{(6.3)^k}{k!}$ for $k = 0, 1, 2, \dots$

b. (4 points) $f(x) = \begin{cases} cx^{5.3}(1-x)^{7.14} & \text{for } x \in (0, 1), \\ 0 & \text{otherwise.} \end{cases}$

c. (4 points) $f(k) = c \binom{40}{k} \binom{50}{13-k}$ for $k = 0, 1, 2, \dots, 13$.

d. (4 points) $f(x) = \begin{cases} ce^{-5.3x+2} & \text{for } x > 0, \\ 0 & \text{otherwise.} \end{cases}$

e. (4 points) $f(x) = ce^{-5.3x^2+2}$ for $x \in \mathbb{R}$.

Problem 2. (20 points) Let $X = \text{Unif}(0, 1)$ be uniform on $[0, 1]$.

a. (5 points) For which parameters (α, β) is $X = \text{Unif}(0, 1)$ the same as the Beta distribution with parameters (α, β) ?

b. (15 points) Given $\gamma > 0$ a positive real number, and $X = \text{Unif}(0, 1)$ as before, prove that $Y = X^\gamma$ is Beta distributed with parameters $\alpha = \frac{1}{\gamma}, \beta = 1$, that is, $Y = \text{Beta}(\frac{1}{\gamma}, 1)$.

Problem 4. (20 points) Let (X_1, X_2) have a bivariate normal distribution with means $\mathbf{E}X_1 = 10$, $\mathbf{E}X_2 = 8$, standard deviations $\sigma(X_1) = 1$, $\sigma(X_2) = 3$, and correlation $\rho(X_1, X_2) = 0.5$. Let $Y = X_1 - X_2$.

a. (5 points) What choice of an estimate m_0 for Y will minimize the mean square error $\mathbf{E}((Y - m_0)^2)$?

b. (5 points) What choice of an estimate m_1 for Y will minimize the mean absolute error $\mathbf{E}(|Y - m_1|)$?

c. (5 points) Choosing m_0 as in part (a), compute $\mathbf{E}((Y - m_0)^2)$.

d. (5 points) Express $\mathbf{Pr}(X_1 > X_2)$ in the terms of the cdf $\Phi(z) = \int_{-\infty}^z e^{-\frac{t^2}{2}} dt$ for a standard normal $Z = \mathcal{N}(0, 1)$ having mean 0, standard deviation 1.

Problem 5. (20 points) Let X be a Poisson random variable with unknown mean λ , and you have been told, *a priori*, that λ comes from a Gamma distribution $\text{Gamma}(2, 3)$ with parameters $\alpha = 2, \beta = 3$.

You sample X and obtain the value $X = 5$.

Show that the *posterior* distribution for λ still follows a Gamma distribution $\text{Gamma}(\hat{\alpha}, \hat{\beta})$, and find the new values $(\hat{\alpha}, \hat{\beta})$ explicitly.