

Name: _____

Signature: _____

Math 5651 Lecture 002 (V. Reiner) Midterm Exam I
Thursday, February 25, 2016

This is a 115 minute exam. No books, notes, calculators, cell phones, watches or other electronic devices are allowed. You can leave answers as fractions, with binomial or multinomial coefficients unevaluated.

There are a total of 100 points. To get full credit for a problem you must show the details of your work. Answers unsupported by an argument will get little credit. Do all of your calculations on this test paper.

Problem	Score
1.	_____
2.	_____
3.	_____
4.	_____
5.	_____
6.	_____
7.	_____
Total:	_____

Reminders:

$$\Pr(A_1 \cup \dots \cup A_n) = \sum_{k=1}^n (-1)^{k-1} \sum_{1 \leq i_1 < \dots < i_k \leq n} \Pr(A_{i_1} \cap \dots \cap A_{i_k})$$

$$S = B_1 \sqcup \dots \sqcup B_n \Rightarrow \Pr(A) = \sum_{i=1}^n \Pr(A \cap B_i) = \sum_{i=1}^n \Pr(A|B_i)\Pr(B_i)$$

and Bayes Theorem $\Pr(B_i|A) = \Pr(A|B_i)\Pr(B_i)/\Pr(A)$

$$\mathbf{E}X = \sum_k k \cdot f(k) \quad \text{for a discrete random variable with p.f. } f(k)$$

$$X = \text{Bin}(n, p) \text{ has p.f. } f(k) = \binom{n}{k} p^k (1-p)^{n-k} \text{ for } k \in \{0, 1, 2, \dots, n\}$$

$$X = \text{Hypergeom}(A, B, n) \text{ has p.f. } f(k) = \binom{A}{k} \binom{B}{n-k} / \binom{A+B}{n} \text{ for } k \in \{0, 1, 2, \dots, \min\{A, n\}\}$$

$$X = \text{Poi}(\lambda) \text{ has p.f. } f(k) = e^{-\lambda} \frac{\lambda^k}{k!} \text{ for } k \in \{0, 1, 2, \dots\}$$

Problem 1. (*10 points*) When rolling a fair 6-sided dice having 1, 2, 3, 4, 5, 6 on its sides n times, consider the two events A, B where A is rolling a number 3 or less every time, and B is rolling the same number every time. Are A and B dependent or independent? You must support your answer with calculations to receive any credit.

Problem 2. (*15 points*) If I choose a rearrangement of the 11 letters in the word "MISSISSIPPI" into a possibly nonsensical string of 11 letters, with all rearrangements equally likely, then what is the probability that *at least* one (and possibly more than one) of the following three events occurs?:

- The four letters "SSSS" appear all adjacent.
- The four letters "IIII" appear all adjacent.
- The two letters "PP" appear adjacent.

Problem 3. (15 points) Your friend Bill Shakespeare tells you that he has written a new play, either a comedy or a tragedy. However, you know that whenever he says this, it is *always* written by someone else, a ghost-writer which is either Chris Marlowe, Frank Bacon, or Benny Jonson, each with their own fixed frequencies for writing tragedies versus comedies:

	Marlowe	Bacon	Jonson
fraction of time used as ghost-writer	1/2	1/4	1/4
% chance of writing a tragedy	90%	50%	20%

a. (5 points) Before he shows you the play, what is the probability that it is a comedy?

b. (10 points) You read it, and it is a comedy. What is the probability that Jonson wrote it?

Problem 4. (*15 points total*) A group of 14 women and 8 men people pair off to make 11 pairs of swimming buddies. If all possible pairings equally likely, what is the probability that all the pairs are single-sex, that is, woman-woman or man-man?

Problem 5. (*15 points total*) Prove that if the events A, B, C are (jointly/mutually, not just pairwise) independent, then the events A^c and $B \cup C$ are also independent.

Problem 6. (*15 points total*) You have two coins in your pocket, one *fair* coin having probability of heads $1/2$, and one *unfair* coin in which the probability of heads is $2/3$. You pull one of the two out of your pocket, with either coin equally probable, and start flipping it, generating a sequence of heads and tails.

a. (5 points) What is the probability that in a total of 10 flips you get at most 8 heads?

b. (5 points) What is the probability that the first tails occurs exactly on the 5^{th} flip?

c. (5 points) After you pull it out of your pocket, you do one test flip, and get tails. What is the probability that you pulled the fair coin out of your pocket?

Problem 7. (15 points) Assume $X = Poi(\lambda)$ is a Poisson random variable with mean λ , and let $Y = X^2$, so that Y only takes on the values k^2 for $k = 0, 1, 2, \dots$, and

$$\Pr(Y = k^2) = e^{-\lambda} \frac{\lambda^k}{k!}.$$

What is the expected value $\mathbf{E}(Y)$ of Y ?

(Hint: I think it helps to rewrite $k^2 = k(k - 1) + k$.)