

A formula for F -Polynomials in terms of C -Vectors and Stabilization of F -Polynomials

Meghal Gupta

September 20, 2018

Abstract

In this paper, we present a general formula for the coefficients of F -polynomials in terms of C -matrix entries. We then demonstrate the effectiveness of the formula by using it to derive simple formulas for specific classes of quivers. Work has been done to do these cases in ad-hoc combinatorial ways, but our formula recovers them with a general method. Secondly, we investigate convergence for F -polynomials. In themselves, they do not converge, but by changing bases using the C -matrix, they conjecturally do. Since our formula relates C -matrix entries to coefficients, we are able to apply it to make considerable progress on the conjecture. In particular, we show stability for green mutation sequences. Finally, we look at exact formulas for these stable deformed F -polynomials in instances where they illustrate properties that the F -polynomials themselves do not.

1 Introduction

Cluster algebras, introduced by Fomin and Zelevinsky in 2002 [4], are commutative algebras with a distinguished set of generators, known as cluster variables. Their original motivation came from an algebraic framework for total positivity and canonical bases in Lie Theory. In recent years, connections have been made to other areas, such as physics, quiver representations, Calabi-Yau categories, and Poisson geometry.

The cluster variables are grouped into sets of constant cardinality v , the **clusters**, and the integer v is called the rank of the cluster algebra. Start with an initial cluster \mathbf{x} (together with a skew symmetrizable integer $v \times v$ matrix B and a coefficient vector $\mathbf{p} = \langle (p_1^+, p_1^-), (p_2^+, p_2^-), \dots, (p_v^+, p_v^-) \rangle$ whose entries are elements of a torsion-free abelian group. We denote the entry in the k th row and i th column of B as $B[k, i]$. The set of cluster variables is obtained by repeated application of so called **mutations**. To perform a mutation at k , replace x_k in the cluster with

$$x'_k = \frac{1}{x_k} \left(p_k^+ \prod_{B[k,i]>0} x_i^{B[k,i]} + p_k^- \prod_{B[k,i]<0} x_i^{-B[k,i]} \right)$$

This creates a new cluster variable. By a result of Fomin and Zelevinsky [4] known as the Laurent phenomenon, this new cluster variable will always be a Laurent polynomial. There is also a corresponding change to the matrix B and vector p , which will be defined properly in a later section. When one sets each initial x_i to 1 and initial p vector as $\langle y_1, \dots, y_v \rangle$, each cluster variable is a polynomial with indeterminates $\langle y_1, \dots, y_v \rangle$. These polynomials are called F -polynomials. It turns out that the coefficients of monomials in F -polynomials are the same as those we would have gotten by setting each initial y_i to 1 and leaving the initial x vector as indeterminates $\langle x_1, \dots, x_v \rangle$. As such, results about coefficients of F -polynomials also apply to cluster variables.

F -Polynomials have been an interesting area of study since the inception of the field. Fomin and Zelevinsky conjectured in [4] that every F -polynomial has all positive coefficients. This conjecture was later proven in the skew-symmetric case in [11] and then the entire skew-symmetrizable case in [7]. Nonetheless, we are

far from completely understanding them. For one, not very many explicit formulas exist for the coefficients of F -polynomials. Since they are defined through recurrence relations, it is not clear that we can actually get a closed form formulation instead. The main result of our paper is to derive a closed form based only on the C -matrix entries, which are related to the p -vectors defined above. Some of the most general previous results regarding F -polynomial explicit forms are the following:

- A formula in terms of the Euler-Poincare characteristic of quiver Grassmannians obtained in [2].
- A combinatorial interpretation for cluster algebras from surfaces given in [14].
- A formula for cluster variables corresponding to string modules as a product of 2×2 matrices obtained in [1].
- A formula using broken lines and reflections in [7].

The last result is the only one that is completely general; the others are restrictive in the cluster algebras they account for. The first is mostly general, restricting only in that it only applies to skew-symmetric cluster algebras. However, the first and last have limited computational power, as the coefficients involve objects that are hard to understand and compute.

As said earlier, the main result of this paper is to present a formula for the F -polynomial that is completely general and completely explicit. This is in itself interesting, because it shows that the recurrence relation for F -polynomials can be solved out. Moreover, we show that our formula is useful. We demonstrate that this formula has computational power, meaning its terms are easy to understand, unlike the formula using the Euler-Poincare characteristic of quiver Grassmannians. We demonstrate this by using it to easily obtain algebraic formulas for F -polynomials of specific classes of quivers, that are as simple or simpler than previously known. Though our formula is general, our examples and applications of it restrict to the skew-symmetric case, because that is where patterns are most evident. Furthermore, the formula has an application to geometry in this case, where the cluster algebra can be interpreted with quivers. The formula for the F -polynomial in terms of the Euler-Poincare characteristic of quiver Grassmannians mentioned earlier in [2] says the following:

$$F_n = \sum_e \chi(Gr_e(\phi_n)) \prod_{i=1}^v y_i^{e_i}$$

Here, e is a vector in \mathbb{N}^v and ϕ_n is a representation of the original quiver defined by certain mutation rules. As such, our expression for the coefficient of a given monomial in an F -Polynomial also gives a formula for the corresponding Euler-Poincare characteristic of quiver Grassmannians associated to dimension vectors parameterized by e . This is interesting, because these are often very difficult to understand and compute. Finally, it is well known that F -polynomial coefficients for specific quivers represent various combinatorial objects; for example pyramid partitions [3], paths in surfaces with marked points [17], and snake graphs [15]. Therefore, our F -polynomial formula also provides a way to count these combinatorial objects.

Secondly, we use our formula to make progress on a conjecture made by Eager and Franco in Section 9.5 [3]. They explore convergence of F -polynomials in the skew symmetric case. In themselves, F -polynomials do not converge, but when one changes basis; i.e. acts with an appropriate linear operator on the exponent vector of each monomial, they appear to converge. This phenomenon and precise linear operator was conjectured in [3] and formalized by Grace Zhang in [18]. This bears a striking resemblance to our formula for the coefficients of F -Polynomials, and we use our results to prove Eager and Franco's conjecture for a certain class of quivers known as green quivers. This includes the quiver they originally conjectured stabilization for, the $dP1$ quiver, known in combinatorics for giving rise to the Somos-4 sequence. This also extends results [18], which prove stabilization for some specific green quivers.

2 Definitions

We start with the general definitions, and later state the simplified definitions for the skew-symmetric case, since that is the case we study most closely. The way we define F -polynomials is purely combinatorial; we

do not use the cluster algebra notation directly. In both cases, we would start with a matrix B and mutation sequence v_1, \dots, v_n to obtain the F -polynomial F_n . In our definitions, the matrix B is essentially replaced by the notion of a generalized quiver defined below in Section 2.1. An interested reader should look at [4] and [5] to understand why the cluster algebra definition of an F -polynomial is the same.

2.1 General Definitions

Definition 2.1 (Generalized Quiver). A **generalized quiver** Q is a directed graph on vertices $1, \dots, |Q|$, where every edge is associated to one of its incident vertices, along with a Laurent polynomial label at every vertex. However the edges must satisfy certain properties:

- There are no 2-cycles involving edges both associated to the same vertex.
- When written in a matrix B with entry $B[i, j]$ being the number of edges associated to j going from $j \rightarrow i$ (negative if edges go in), B is skew-symmetrizable.

Intuitively, a generalized quiver is a generalized version of a quiver, reflecting the fact that the “adjacency matrix” B need not be skew-symmetric.

Definition 2.2 (Generalized Quiver Mutation). An **generalized quiver mutation** at vertex i is defined by changing both the quiver around vertex i and the Laurent polynomial at vertex i . The quiver changes are defined as follows.

- For every path $j \rightarrow i \rightarrow k$ with the first edge associated to i and the second to k , draw a new edge from j to k associated to k .
- For every path $j \rightarrow i \rightarrow k$ the first edge associated to j and the second to i , draw a new edge from j to k associated to k . Reverse the direction of every edge pointing with i as a vertex.
- Delete all 2-cycles involving edges both associated to the same vertex.

The Laurent polynomial is replaced as follows. Let V_k denote the Laurent polynomial currently on vertex k . Replace V_i with the following function P .

$$P = \frac{\prod_{j \in S_1} V_j + \prod_{j \in S_2} V_j}{V_i}$$

where S_1 is the set of vertices with multiplicity pointing into i associated to i and S_2 is the set of vertices with multiplicity of edges pointing out of i associated to i .

Notice that the vertex i may no longer have a Laurent polynomial label. However, the following theorem by Fomin and Zelevinsky rectifies that.

Theorem 2.3 (Laurent Phenomenon [4]). *Take a generalized quiver Q with starting polynomials x_1, \dots, x_v at each corresponding vertex. For any mutation sequence, the function P as defined above is actually a Laurent polynomial.*

Definition 2.4 (Framed Generalized Quiver). The **framed generalized quiver** of a generalized quiver Q with labels (polynomials) 1 at every vertex is the quiver formed by adding a vertex i' for every vertex i with an edge pointing from i' to i associated to i . The new vertices i' have the label y_i . A **base vertex** of the framed generalized quiver is a vertex that is also a vertex of the original quiver.

Definition 2.5 (F -Polynomial). The n th F -Polynomial of a framed generalized quiver Q accompanied with a mutation sequence v_1, \dots, v_n of vertices is the polynomial at vertex v_n after performing mutations at v_1, \dots, v_n sequentially.

Definition 2.6 ($V_n(i)$ or V_i). Given a quiver Q and a sequence of mutations v_1, \dots, v_n , $V_n(i)$ is the F -polynomial on vertex i after the n mutations. We will usually denote this F -polynomial more simply as V_i if we are otherwise clear about the termination point n for the sequence of mutations.

Definition 2.7 (C -matrix). The **C -matrix** of the framed generalized quiver Q and mutation sequence v_1, \dots, v_n with $|Q|$ base vertices is the $|Q| \times |Q|$ matrix with entry (i, j) as the number of edges from frozen vertex i' to base vertex j associated to j . It is denoted C_n .

Definition 2.8 (A_i). Given a framed generalized quiver Q and a sequence of mutations v_1, \dots, v_i , let A_i^g be the identity matrix with entry (v_i, v_i) replaced with -1 and for every arrow pointing from v_i to another base quiver vertex v associated to v , add 1 to the entry (v_i, v_v) . A_i^r is defined analogously with arrows pointing into v_i . We will just refer to the relevant A_i^g or A_i^r by A_i , where the relevant one is what corresponds to the color of r_i as will be defined soon. The other one will be A_i^* .

Lemma 2.9 (C -Matrix Facts, Theorem 5.6 [10]). *Given a framed quiver Q and a sequence of mutations v_1, \dots, v_n , $C_n = A_1 A_2 \cdots A_n$, where for each i you choose g if all the arrows are pointing into v_i before the i th mutation, and r otherwise. Also, C_n is invertible.*

Definition 2.10 (E_i). Given a framed generalized quiver Q and a sequence of mutations v_1, \dots, v_i , let E_i^g be the identity matrix with entry (v_i, v_i) replaced with -1 and for every arrow pointing from v_i to another base quiver vertex v associated to v_i , add 1 to the entry (v_i, v_v) . E_i^r is defined analogously with arrows pointing into v_i . We will just refer to the relevant E_i^g or E_i^r by E_i , where the relevant one is what corresponds to the color of r_i as will be defined soon. The other one will be E_i^* .

Definition 2.11 (D -Matrix). The **D -matrix** is similar to the C -matrix, defined slightly differently though: $D_n = E_1 \cdots E_n$. Note that D_n is also invertible. The D -matrix between two steps is defined as follows. For two indices m and n , let $D_{m,n}$ be $D_n D_m^{-1}$. By extension, $D_n = D_{0,n}$.

Definition 2.12 (r_i). Given the quiver formed by a framed quiver Q and a mutation sequence v_1, \dots, v_i , let r_i be the monomial that is the product of the frozen nodes pointing into/from vertex v_i . The r_i is called green if the frozen nodes are pointing in before the final mutation (out afterwards) and red if they are all pointing out.

Theorem 2.13 (Sign Coherence [7]). *Given the quiver formed by a framed quiver Q and a mutation sequence v_1, \dots, v_i , all the frozen vertices pointing into any vertex v at the end of the process are in the same direction (all in or all out), and there will be at least one arrow. Thus, every vertex will be green or red.*

We call a mutation sequence green if all the mutations are green. We often just say the quiver is green if it is clear what mutation sequence we are referring to.

The next two definitions bare no obvious significance, but simplify expressions in future sections. Loosely, the particular values $a_{i,j}$ and $b_{i,j}$ are important because they are the building blocks of coefficients of monomials in the F -polynomials.

Definition 2.14 ($a_{i,j}$ and $b_{i,j}$). Given a framed quiver Q and a mutation sequence v_1, v_2, \dots , for two indices $i \leq j$, let $a_{i,j} = D_{i,j}^{-1}[v_j, v_i]$. Similarly, let $b_{i,j} = E_j^* E_j D_{i,j}^{-1}[v_j, v_i]$, or equal to 0 when $i = j$. Let $\delta_{i,j}$ be 1 if r_i and r_j are the same color, and -1 otherwise. Mutation 0 counts as red for the purposes of assigning $\delta_{0,i}$.

Definition 2.15 (W). Given a framed quiver Q and a mutation sequence v_1, \dots, v_n , and a sequence of indices w_1, \dots, w_k where the w_i 's are all between 1 and n inclusive, let

$$W(n, w_1, \dots, w_k) = \prod_{i=1}^k \left((a_{w_i, n}) + \sum_{j=i+1}^k (-a_{w_i, w_j} + b_{w_i, w_j}) \right).$$

Definition 2.16 (ϕ). We define $\phi(\mathbf{w}_1, \dots, \mathbf{w}_k)$ to be the number of distinct permutations of w_1, \dots, w_k divided by $k!$. Note that this is just $\frac{1}{\prod a_i!}$ where the a_i are the number of occurrences of each of the distinct values taken on by w_1, \dots, w_k .

2.2 Simplifications in the Skew-symmetric Case

In the case where B is skew-symmetric, for any edge between two base vertices, the number of edges associated to each vertex is the same. As such, we can define the quiver, rather than generalized quiver, so that only one set of edges is included, and association can be disregarded. The analogous definitions are straightforward, but we redefine for completeness. To understand that all proofs completed in the general case also apply here, one should check that all the variables defined, C , A , $a_{i,j}$, $b_{i,j}$, r_i , etc. are all the same when returning the quiver to its generalized quiver form.

Definition 2.17 (Quiver). A **quiver** Q is a directed graph, along with a Laurent polynomial label at every vertex. $|Q|$ denotes the number of vertices in Q . The directed graph must have no 2-cycles.

Any quiver can be viewed, instead, as a generalized quiver by replacing each edge with two edges, one associated to each incident vertex.

Definition 2.18 (Quiver Mutation). An **quiver mutation** at vertex i is defined by changing both the quiver around vertex i and the Laurent polynomial at vertex i . The quiver changes are defined as follows.

- For every path $j \rightarrow i \rightarrow k$, draw a new edge from j to k .
- For every path $j \rightarrow i \rightarrow k$, draw a new edge from j to k . Reverse the direction of every edge pointing with i as a vertex.
- Delete all 2-cycles.

The Laurent polynomial is replaced as follows. Let V_k denote the Laurent polynomial currently on vertex k . Replace V_i with the following polynomial P .

$$P = \frac{\prod_{j \in S_1} V_j + \prod_{j \in S_2} V_j}{V_i}$$

where S_1 is the multiset of edges pointing into i associated to i and S_2 is the multiset of edges pointing out of i .

Definition 2.19 (Framed Quiver). The **framed quiver** of a quiver Q with labels 1 at every vertex is the quiver formed by adding a vertex i' for every vertex i with an edge pointing from i' to i . The new vertices i' have the label y_i . A **base vertex** of the framed quiver is a vertex that is also a vertex of the original quiver.

Definition 2.20 (C -matrix). The **C -matrix** of the quiver formed by a framed quiver Q and mutation sequence v_1, \dots, v_n with $|Q|$ base vertices is the $|Q| \times |Q|$ matrix with entry (i, j) as the number of edges from frozen vertex i' to base vertex j . It is denoted C_n . The C -matrix between two steps is defined as follows. For two indices m and n , let $C_{m,n}$ be $C_n C_m^{-1}$. It is worth noting that $C_n = C_{0,n}$.

Definition 2.21 (A_i). Given a framed quiver Q and a sequence of mutations v_1, \dots, v_i , let A_i^g be the identity matrix with entry (v_i, v_i) replaced with -1 and for every arrow pointing from v_i to another base quiver vertex v , add 1 to the entry (v_i, v) . A_i^r is defined analogously with arrows pointing into v_i . We will just refer to the relevant A_i^g or A_i^r by A_i , where the relevant one is what corresponds to the color of r_i as will be defined soon. The other one will be A_i^* .

The A -matrix is the same as the E -Matrix. Both will be denoted by A . The C -matrix is the same as the D -matrix. Both will be denoted by C . Also note the following:

Definition 2.22 ($a_{i,j}$ and $b_{i,j}$). Given a framed quiver Q and a mutation sequence v_1, v_2, \dots , for two indices $i \leq j$, let $a_{i,j} = C_{i,j}^{-1}[v_j, v_i]$. Similarly, let $b_{i,j} = A_j^* A_j C_{i,j}^{-1}[v_j, v_i]$, or equal to 0 when $i = j$. Let $\delta_{i,j}$ be 1 if r_i and r_j are the same color, and -1 otherwise. Mutation 0 counts as red for the purposes of assigning $\delta_{0,i}$.

Definition 2.23 (W). Given a framed quiver Q and a mutation sequence v_1, \dots, v_n , and a sequence of indices w_1, \dots, w_k where the w_i 's are all between 1 and n inclusive, let

$$W(n, w_1, \dots, w_k) = \prod_{i=1}^k \left((a_{w_i, n}) + \sum_{j=i+1}^k (-a_{w_i, w_j} + b_{w_i, w_j}) \right).$$

We will use $a_{i,j}$, $b_{i,j}$ and W regardless of whether we are dealing with a quiver or generalized quiver.

2.3 Definitions for Stabilization

Definition 2.24 (Matrix Action on Polynomial). Given an $v \times v$ matrix M and a polynomial P in x_1, \dots, x_v , we let $M(P)$ denote the polynomial formed when M acts on the vector of exponents for each monomial in P .

Definition 2.25 (Deformed F -Polynomial). The n th S -polynomial (deformed F -polynomial) of a framed quiver Q with a mutation sequence v_1, \dots, v_n is defined as $-C^{-1}(F_n)$.

Definition 2.26 (Fundamental r_i). An r_i of a framed quiver Q and mutation sequence v_1, v_2, \dots is called **fundamental** if it is not expressible as the product of at least 2 other r_j 's with $j < i$. Let the set M_n denote the set of fundamental r_i for a framed quiver Q and sequence of mutations v_1, \dots, v_n . M is the set of fundamental r_i for the entire mutation sequence v_1, v_2, \dots .

Definition 2.27 (Basic). Given a framed quiver Q and a mutation sequence v_1, \dots, v_n , a monomial m is called **basic** if its coefficient is nonzero in a polynomial that appears on one of the vertices after n mutations. In other words, its coefficient is nonzero in some element of the cluster. Let P_n denote the set of basic monomials after n mutations.

3 F -Polynomial Formula

3.1 Derivation

Theorem 3.1. Given a framed quiver or a framed generalized quiver Q and a mutation sequence v_1, \dots, v_n , F_n can be calculated as follows. Let W be the set of sequences \mathbf{w} with $1 \leq w_1 \leq \dots \leq w_k \leq n$.

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) W(n, w_1, \dots, w_k) \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{0, w_j} C_{0, w_j} [i, v_{w_j}]} \quad (1)$$

Consequently, the coefficient of a given monomial p is the sum of all sequences $1 \leq w_1 \leq \dots \leq w_k \leq n$ so that

$$p = \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{0, w_j} C_{0, w_j} [i, v_{w_j}]}$$

of

$$\phi(w_1, \dots, w_k) W(n, w_1, \dots, w_k).$$

Proof. It suffices to prove the statement in the case of framed generalized quivers.

Consider the steps v_1, \dots, v_n of the mutation process on the quiver Q that results in F_n . At each step i , define \succ_i and \prec_i as follows. If r_i is green, any edge in the quiver before the i th mutation associated to a denoted $a \rightarrow b$ is now denoted $a \succ_i b$. Similarly, $a \leftarrow b$ is denoted $a \prec_i b$. If r_i is red, the reverse occurs: \succ_i replaces \leftarrow , and \prec_i replaces \rightarrow . Thus, \succ replaces \rightarrow or \leftarrow corresponding to the direction of the frozen vertices into v_i . At each step i , the following occurs to calculate F_i . We have

$$F_i V_{v_i} = \prod_{v_i \succ j} V_j + r_i \prod_{v_i \prec j} V_j.$$

We modify this mutation process: instead of plugging in the monomial r_i , replace r_i with the variable z_i . Define the analog of F_i as Z_i and the analog of V_j as ZV_j , so ZV_j is the Z polynomial at the j th vertex. Thus, Z_n will be some rational function in the z_i 's, computed by the following recurrence relation:

$$Z_i ZV_{v_i} = \prod_{v_i \succ j} ZV_j + z_i \prod_{v_i \prec j} ZV_j$$

and $\dots, F_{-1}, F_0 = 1$. When we plug in r_i for each z_i , we get back F_n . As such, to get the coefficient of a given monomial p , do the following: express Z_n in its MacLaurin series form, i.e. an element of $\mathbb{C}[[z_1, \dots, z_n]]$. Then, add the coefficients over all sequences w_1, \dots, w_k so that $p = r_{w_1} \cdots r_{w_k}$.

Now, we compute the coefficient of a given $z_{w_1} \cdots z_{w_k}$. We show it is equal to

$$\phi(w_1, \dots, w_k) W(n, w_1, \dots, w_k).$$

Encode the sequence \mathbf{w} instead as two sequences \mathbf{q} and \mathbf{m} where \mathbf{q} consists of the distinct terms of \mathbf{w} in increasing order, and \mathbf{m} is the corresponding multiplicity of each. Let $l = |\mathbf{q}|$. The coefficient of the term

$$z_{w_1} \cdots z_{w_k} = z_{q_1}^{m_1} \cdots z_{q_l}^{m_l}$$

in Z_n is the same as the one in the polynomial computed as follows: look at the entire modified mutation process, except on a mutation step i not corresponding to a q_j , replace z_i with 0. This is because replacing all irrelevant z_i 's with 0 does not affect the coefficient of the desired term. Call this new polynomial Y_n . YV_{v_i} is defined analogously as well. Consider the following polynomials, L_1, \dots, L_l .

$$\begin{aligned} L_1 &= (1 + z_{q_1}) \\ L_2 &= (1 + z_{q_2}(1 + z_{q_1})^{-a_{q_1, q_2} + b_{q_1, q_2}}) \\ &\vdots \\ &\vdots \\ &\vdots \\ L_l &= \left(1 + z_{q_l} L_{l-1}^{-a_{q_{l-1}, q_l} + b_{q_{l-1}, q_l}} \cdots L_1^{-a_{q_1, q_l} + b_{q_1, q_l}}\right) \end{aligned} \tag{2}$$

Let

$$H_i := \prod_{j=1}^r L_j^{a_{q_j, i}}.$$

where r is the final q_r preceding i (so H_n would be $\prod_{j=1}^l L_j^{a_{q_j, i}}$). We show that at any given step i , $Y_i = H_i$. We do so by induction.

Base Case: $i \leq q_1$: This case is clear. We have

$$Y_{q_1} = H_{q_1} = (1 + z_{q_1})$$

and for $i < q_1$, we have

$$Y_i = H_i = 1.$$

We have the following recurrence relation.

$$Y_i YV_{v_i} = \prod_{v_i \succ j} YV_j + z_i \prod_{v_i \prec j} YV_j$$

I claim that the relation

$$H_i HV_{v_i} = \prod_{v_i \succ j} HV_j + z_i \prod_{v_i \prec j} HV_j$$

holds as well. Because of the inductive hypothesis, proving this recurrence relation would suffice to show the polynomials are equal.

Case 1: $i \neq q_r$: We have $z_i = 0$, so we want to show

$$\begin{aligned} H_i H V_{v_i} &= \prod_{v_i \succ j} H V_j + z_i \prod_{v_i \prec j} H V_j \\ \iff H_i &= \frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}}. \end{aligned}$$

We look at the exponent for each L_s in H_i and show that it is, in fact, the same on both sides of the equation. The exponent on the right hand side, denoted $e_s \left(\frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}} \right)$ is calculated as follows:

$$\sum_{v_i \succ j} e_s(H V_j) - e_s(H V_{v_i}).$$

Let i_j be the last index of a mutation at vertex j . By the inductive hypothesis,

$$e_s \left(\frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}} \right) = \sum_{v_i \succ j} a_{q_s, i_j} - a_{q_s, i_{v_i}}.$$

Recall that

$$D_{i,j} = E_{i+1} \cdots E_j \implies D_{i,j}^{-1} = E_j \cdots E_{i+1}$$

since the E_i are involutions. Multiplication by any E_i only changes the corresponding row, so since $a_{i,j} = D_{i,j}^{-1}[j, i]$, we can replace terms of the form a_{q_s, i_j} with $D_{q_s, i-1}^{-1}[j, v_{q_s}]$, since no entry in the row of the corresponding matrix changes. It's useful to note this also stays 0 when L_s corresponds to a term that appears after one of our HV terms is calculated. Thus, we have as our expression:

$$e_s \left(\frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}} \right) = \sum_{v_i \succ j} D_{q_s, i-1}^{-1}[j, v_{q_s}] - D_{q_s, i-1}^{-1}[v_i, v_{q_s}].$$

However, the right hand side is exactly the operation corresponding to obtaining the entry $D_{q_s, i}^{-1}[v_i, v_{q_s}] = a_{q_s, i}$ from how E_i is defined: add the contributions from all the vertices pointing opposite of the frozen vertices, and subtract the current vertex's contribution. This exactly proves our desired statement that

$$e_s(H_i) = e_s \left(\frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}} \right).$$

Case 2: $i = q_r$: Just as in the last case, we can try to compute

$$\frac{\prod_{v_i \succ j} H V_j + z_i \prod_{v_i \prec j} H V_j}{H V_{v_i}}.$$

By the first case,

$$\frac{\prod_{v_i \succ j} H V_j}{H V_{v_i}}$$

is H_i without the contribution from L_r . We now have to show that adding $\frac{z_i \prod_{v_i \succ j} H V_j}{H V_{v_i}}$ gives the same contribution as multiplying by L_r . By the same logic as the proof from the first case, we find the term

$$\frac{\prod_{v_i \prec j} H V_j}{H V_{v_i}}.$$

Again we find e_s of the expression for each k . We get

$$e_s \left(\frac{\prod_{v_i \prec j} H V_j}{H V_{v_i}} \right) = \sum_{j \succ v_i} D_{q_s, i-1}^{-1}[j, v_{q_s}] - D_{q_s, i-1}^{-1}[v_i, v_{q_s}].$$

This is exactly the operation of E_i^* , so we get

$$e_s \left(\frac{\prod_{j \succ v_i} HV_j}{HV_{v_i}} \right) = b_{q_j, i}.$$

Now that we have these identities, we can show that our recurrence relation in fact gives H_i .

$$\begin{aligned} & \frac{\prod_{v_i \succ j} HV_j + z_i \prod_{v_i \prec j} HV_j}{HV_{v_i}} \\ &= \prod_{j=1}^{r-1} L_j^{a_{q_j, i}} + z_i \prod_{j=1}^{r-1} L_j^{b_{q_j, i}} \\ &= (1 + z_i \prod_{j=1}^{r-1} L_j^{b_{q_j, i} - a_{q_j, i}}) \prod_{j=1}^{r-1} L_j^{a_{q_j, i}} \\ &= L_r \prod_{j=1}^{r-1} L_j^{a_{q_j, i}} \\ &= H_i \end{aligned}$$

which is what we desired.

Now we must actually compute the coefficient of $z_{w_1} \cdots z_{w_k}$ in $H_n = \prod_{j=1}^k L_j^{a_{q_j, n}}$. We prove that it is $\phi(w_1, \dots, w_k) W(n, w_1, \dots, w_k)$ by induction on $l = |\mathbf{q}|$.

Base Case: $|\mathbf{q}| = 1$: Then $H_n = L_1^{a_{q_1, n}}$. By the extended binomial theorem, the coefficient of $z_{q_1}^{m_1}$ is in fact

$$\frac{(a_{q_1, i})(a_{q_1, i} - 1) \cdots (a_{q_1, i} - m_1 + 1)}{m_1!}.$$

Inductive step: $|\mathbf{q}| = l$: If we instead compute the coefficient of the term $q_2^{m_2} \cdots q_l^{m_l}$, then by the inductive hypothesis we get the desired result. The corresponding L_i , which we call L'_i , we have are defined as follows.

$$\begin{aligned} L'_2 &= (1 + z_{q_2}) \\ L'_3 &= (1 + z_{q_3}(1 + z_{q_2})^{-a_{q_1, q_2} + b_{q_1, q_2}}) \\ &\vdots \\ &\vdots \\ &\vdots \\ L'_l &= \left((1 + z_{q_l})(L'_{l-1})^{-a_{q_{k-1}, q_k} + b_{q_{k-1}, q_k}} \cdots L'_1^{-a_{q_1, q_k} + b_{q_1, q_k}} \right) \end{aligned} \tag{3}$$

This gives $H'_n = \prod_{j=1}^k L'_j^{a_{q_j, n}}$. By inspecting the way the L_i 's are formed, we see that we can form H_n by replacing every instance of z_{q_i} with $z_{q_i}(1 + z_{q_1})^{-a_{q_1, q_i} + b_{q_1, q_i}}$, and multiplying the entire expression by $(1 + z_{q_1})^{a_{q_1, n}}$. If we expand only q_2, \dots, q_l , leaving the $(1 + z_{q_1})$ expressions, which corresponds to just expanding H'_n , the term

$$z_{q_2}^{m_2} \cdots z_{q_l}^{m_l}$$

corresponds to the term

$$(1 + z_{q_1})^{a_{q_1, n} + \sum_{i=2}^l -m_i a_{q_1, q_i} + m_i b_{q_1, q_i}} z_{q_2}^{m_2} \cdots z_{q_l}^{m_l}.$$

Any other terms cannot have a monomial that has the correct number of factors of all of z_{q_2}, \dots, z_{q_l} . Thus, we only need to find the coefficient of $z_{w_1} \cdots z_{w_k}$ in this monomial. The corresponding term in H'_n already

has coefficient $\phi(w_{m_1+1}, w_k)W(n, w_{m_1+1}, \dots, w_k)$, so the monomial itself is that times

$$(1 + z_{q_1})^{a_{q_1, n} + \sum_{i=2}^l -m_i a_{q_1, q_i} + m_i b_{q_1, q_i}} z_{q_2}^{m_2} \dots z_{q_l}^{m_l}.$$

To find the total coefficient, we multiply this by the coefficient of $z_{q_1}^{m_1}$ in the expansion of

$$(1 + z_{q_1})^{a_{q_1, n} + \sum_{i=2}^l -m_i a_{q_1, q_i} + m_i b_{q_1, q_i}}$$

which by extended binomial theorem is

$$\frac{a_{q_1, n} + \sum_{i=2}^l -a_{q_1, q_i} + b_{q_1, q_i} \dots a_{q_1, n} + \sum_{i=2}^l -m_i a_{q_1, q_i} + m_i b_{q_1, q_i} - m_1 + 1}{m_1!}.$$

Since $a_{i, i}$ is 1 and $b_{i, i}$ is 0, this can be rewritten as

$$\frac{\prod_{j=1}^{m_1} \left(a_{w_1, n} + \sum_{i=m_1+1}^k -a_{w_j, w_i} + b_{w_j, w_i} \right)}{m_1!}.$$

This, in turn, is exactly $\phi(w_1, \dots, w_k)W(n, w_1, \dots, w_k)$, thus proving our inductive step. \square

3.2 Formula Discussion

In this section, we discuss the formula, its terms, and implications.

The formula has the unfortunate property that it is summing over an infinite set of sequences. However, this is easily remedied; it is fairly simple to find an upper bound on the degree of F_n . For example, take the maximum number of edges in the quiver at any stage m and raise this value to the n th power. This must be an upper bound, because at each step the maximum degree of a label on the quiver must be less than m times the previous maximum value. Then, simply sum only over sequences where the expression

$$\prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{0, w_j} C_{0, w_j} [i, v_{w_j}]}$$

has degree less than our upper bound. This is clearly finite, since each $C_{0, w_i} [j, v_{w_i}]$ is positive. Consequently, this at least limits us to only checking sequences with length less than the upper bound on the degree. In the next few sections, we apply our formula to get simpler formulas for specific cases where we know what the relevant entries of the C -matrix are. In many of the cases, we do not limit the sequences we are summing over, if there isn't a specifically interesting way to do so. It is therefore valuable to keep in mind that this procedure would work.

Secondly, let us understand what the C -matrix entries mean. Recall that $C_n [i, j]$ is the number of arrows pointing into vertex j from vertex i' after n mutation steps on the mutation sequence used to obtain C_n . Similarly, $C_{m, n} [i, j]$ is the number of arrows pointing into vertex j from vertex i' after $n - m$ mutation steps where the initial base quiver is actually the original quiver after m mutation steps. There is a caveat though: the color sequence must be the same as it was on the original quiver. If not, we can force it to be the same, loosely by turning a red mutation into a negative analog of a green mutation and vice versa. $C_{m, n} [i, j]$ also makes sense for $n < m$ (simply do the mutations from j to i in reverse), so this lets us understand what the entries of inverse matrices mean, since $C_{m, n} [i, j] = C_{m, n}^{-1} [j, i]$

Another thing to understand is the following. The formula doesn't do a good job of highlighting which sequences of r_i 's are going to contribute negatively and positively to the coefficient of a given monomial. There might be a way to group the sequences, so each group of sequences will be a nonnegative contributor. It would be very cool if we could recover positivity using this formula. It could also help simplify the formula, by reducing the set of sequences we have to sum over, if we know certain groups give 0.

Now we state the analog of the F -polynomial formula for deformed F -polynomials.

Corollary 3.2. *Given a framed quiver Q and a mutation sequence v_1, \dots, v_n , we have that S_n can be calculated as follows. Let W be the set of sequences \mathbf{w} with $0 \leq w_1 \leq \dots \leq w_k \leq n-1$.*

$$S_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) W(n, n - w_k, \dots, n - w_1) \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{n-w_j, n} C_{n-w_j, n}^{-1} [i, v_{n-w_j}]} \quad (4)$$

Consequently, for a given monomial p , over all sequences $0 \leq w_1 \leq \dots \leq w_k \leq n-1$ so that

$$p = \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{n-w_j, n} C_{n-w_j, n}^{-1} [i, v_{n-w_j}]}$$

the value of the coefficient is the sum over all the sequences of

$$\phi(w_1, \dots, w_k) W(n, n - w_k, \dots, n - w_1).$$

Proof. This is Theorem 3.1, with the y_i exponents transformed according to the stabilization rule. \square

This is interesting when we consider the full form of the formula, namely

$$S_n = \sum_{0 \leq w_1 \dots w_k \leq n-1} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((a_{n-w_i, n}) + \sum_{j=1}^{i-1} (-C_{n-w_j, n-w_i}^{-1} [v_{n-w_i}, v_{n-w_j}]) + A_{n-w_i}^* A_{n-w_i} C_{n-w_j, n-w_i}^{-1} [v_{n-w_i}, v_{n-w_j}] \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{n-w_j, n} C_{n-w_j, n}^{-1} [i, v_{n-w_j}]}$$

The terms $C_{n-w_j, n}^{-1} [i, v_{n-w_j}]$ appear both in the exponents of the y_i terms and as a part of the coefficient calculation. This alludes to why C_n^{-1} is a natural operation, and also why it might cause convergence.

4 Exact F -Polynomials

4.1 Symmetric Quivers

We aspire to compute F_n for graphs with useful symmetry properties. We then use this to directly do the next two cases.

Definition 4.1. Let a Symmetric Quiver Q be one with the following properties. Take a quiver Q with base quiver B and the periodic mutation sequence $1, \dots, v, 1, \dots$, with the following properties.

- The base quiver B is reversible; that is reversing all the arrows and relabeling vertex i with $v+1-i$ returns the same quiver.
- The sequence of mutations is cyclic; that is mutating B at 0 returns the original quiver upon relabeling vertex i with $i-1$, where 1 is relabeled with v .
- The base quiver is symmetric; that is swapping i with $v+2-i$ returns a quiver with the same arrows pointing in and out of vertex 1, and fixing vertex 1.
- The quiver is entirely green.

For the quiver Q define

$$f(x_1, \dots, x_v) = -x_1 + \sum_{\text{edge in } B \ 1 \rightarrow i} x_i$$

and

$$g(x_1, \dots, x_v) = -x_1 + \sum_{\text{edge in } B \ i \rightarrow 1} x_i.$$

Define the recurrence relation $s_i = f(s_{i-1}, \dots, s_{i-v})$ and $\dots, s_{-2} = s_{-1} = 0, s_0 = 1$. Let $s'_i = g(s_{i-1}, \dots, s_{i-v})$

Lemma 4.2. Take a symmetric quiver Q and mutation sequence $1, \dots, v, 1, \dots$ with the definitions above. Then,

$$r_k = \prod_{i=1}^v y_i^{s_{k+1-i}}.$$

Proof. We prove the statement by induction.

Base Case: step $k = 1$: We have $r_i = y_1$ and $\prod_{i=1}^v y_i^{s_{1-i}} = y_1$.

Inductive Step: step $k \neq 1$: We are mutating at mutation step k . We need to find the contributors to the frozen vertices pointing into k . Consider the previous cycle of mutation steps, and possible things pointing into or out of vertex k when the vertex was mutated (thus contributing to vertex k) Due to symmetry and cyclicity, the things that were pointing in must be precisely what is pointing out of vertex k now, and vice versa. This easily gives the relation $e_i = f(e_{i-1}, \dots, e_{i-v})$ where e_i is the exponent vector of a given r_i . It is worth verifying that this logic also holds when $k < v$ (vertices that have not yet been mutated will have contribution 0). In turn, the relation proves the desired result for k by induction, and so we have proven the lemma. \square

Theorem 4.3. Take a symmetric quiver Q and mutation sequence $1, \dots, v, 1, \dots$. Let W be the set of sequences $0 \leq w_1 \leq \dots \leq w_k \leq r$. Then

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_j+1-i}}.$$

Proof. We have

$$F_n = \sum_{1 \leq w_1, \dots, w_k \leq n} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((a_{w_i, n}) + \sum_{j=i+1}^k (-a_{w_i, w_j} + b_{w_i, w_j}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k -C_{0, w_j}[i, v_{w_j}]}.$$

Since the quiver is reversible, note that $C_k^{-1} = C_k$ after swapping indices 0 and $v-1$, 1 and $v-2$ etc. Furthermore, $C_{i, k} = C_{k-i}$ after cyclic shifting the indices. With a little work, one can therefore see that $a_{i, k} = a_{k-i}$ and same for b . Further, one can see that $a_{0, i} = s_i$ and $b_{0, i} = s'_i$ because of the lemma. Therefore,

$$F_n = \sum_{1 \leq w_1, \dots, w_k \leq n} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{n-w_i}) + \sum_{j=i+1}^k (-s_{w_j-w_i} + s'_{w_j-w_i}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k -C_{0, w_j}[i, v_{w_j}]}$$


By lemma 4.2, $-C_{0, r}[i, v_r] = s_{k-i}$. This therefore gives

$$F_n = \sum_{1 \leq w_1, \dots, w_k \leq n} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{n-w_i}) + \sum_{j=i+1}^k (-s_{w_j-w_i} + s'_{w_j-w_i}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_j+1-i}}$$

which is the desired expression. \square

4.2 K_r

Definition 4.4. The framed quiver \mathbf{K}_r is the quiver with base quiver having two vertices, 0 and 1, and r arrows from 0 to 1. K_3 is illustrated below.



Definition 4.5. Define the sequence \mathbf{s} as $\dots, s_{-2} = s_{-1} = 0, s_0 = 1$ and $s_{k+2} = ns_{k+1} - s_k$ for $k \geq 0$.

The main focus of the papers [12] and [13] are to find explicit formulas for the F -Polynomials of the quivers K_r . The former derives an algebraic expression that appears to be more complicated than ours. The latter interprets the coefficients as the count of a combinatorial object, and thus is incomparable to our work. It is interesting that an explicit formula follows as a simple corollary of our general formula.

Theorem 4.6. *We can compute F_n for K_r as follows. Let W be the set of sequences \mathbf{w} with $1 \leq w_1 \leq \dots \leq w_k \leq n$.*

$$F_n = 1 + \sum_{\mathbf{w} \in W} \phi(\mathbf{w}) \prod_{i=1}^k \left(s_{n-w_i} - \sum_{j=i+1}^k s_{w_j-w_i} + s_{w_j-w_i-2} \right) y_1^{\sum_{i=1}^k s_{w_i-1}} y_2^{\sum_{i=1}^k s_{w_i-2}}$$

Proof. Notice that K_r is a symmetric quiver. Recall Theorem 4.3 which states that

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_j+1-i}}.$$

Here, \mathbf{s} is defined exactly as in 4.5, and $s'_i = -s'_{i-2}$. Plugging in therefore gives

$$F_n = \sum_{1 \leq w_1 \dots w_k \leq n} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{n-w_i}) - \sum_{j=i+1}^k (s_{w_j-w_i} + s_{w_j-w_i-2}) \right) y_1^{\sum_{j=1}^k s_{w_j-1}} y_2^{\sum_{j=1}^k s_{w_j-2}}$$

This is exactly our desired expression. □

Note that the degree in y_1 of F_n is s_{n-1} , so we only have to sum over sequences with

$$\sum_{i=1}^k s_{w_i-1} \leq s_{n-1}$$

4.3 Gale Robinson Quivers


We now aspire to compute F_n for Gale Robinson Quivers specifically.

Definition 4.7. Define the Gale Robinson Quiver $G_{v,r,t}$ as follows.

- There are v vertices labeled $1, \dots, v$.
- For all $1 \leq i \leq v - r$, draw an arrow $i \rightarrow i + r$, and for all $1 \leq j \leq r$, draw an arrow $j \rightarrow v - r + j$.
- For all $1 \leq i \leq v - t$, draw an arrow $t + i \rightarrow i$, and for all $1 \leq j \leq t$, draw an arrow $v - s + j \rightarrow j$.
- For all $1 \leq i \leq v - r - t$, draw an arrow from $r + i \rightarrow t + i$ and for all $1 \leq j \leq t - r$, draw an arrow $r + j \rightarrow v - t + j$.

- Delete any 2-cycles created in the above process.

We illustrate $G_{7,2,3}$ below.



Lemma 4.8. $G_{v,r,t}$ has the following properties.

- The base quiver B is reversible.
- The sequence of mutations is cyclic.
- The quiver is entirely green.
- Only $1+r$ and $v+1-r$ point out of vertex 1 and $1+t$ and $v+1-t$ point into 0 (this implies it is symmetric).

Proof. The first statement is easy to check, since the arrows are explicitly stated. The second and third are well-known [8]. The fourth, again, is easy to check since the arrows are explicitly stated. \square

Theorem 4.9. Take the quiver $G_{v,r,t}$. Define a sequence \mathbf{s} with s_i as the number of partitions of i into parts $r, v-r$ (0 for negative values of i , 1 for 0). Let W be the set of sequences with $1 \leq w_1 \leq \dots \leq w_k \leq n$. Then

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} - s_{w_j-w_i-v} + s_{w_j-w_i-t} + s_{w_j-w_i-v+t} \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_j-i}}$$

Proof. A Gale Robinson Quiver is in fact a symmetric quiver. Therefore, we can just plug into the formula in the previous section. Define s_i as in the previous section. It is not difficult to see that s_i is the number of partitions of i into parts $r, v-r$ (0 for negative values of i , 1 for 0). We also have $s'_i = s_{i-t} + s_{i-v+t} - s_{i-v}$. Thus, literally plugging in to Theorem 4.3 gives the desired equation of

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} - s_{w_j-w_i-v} + s_{w_j-w_i-t} + s_{w_j-w_i-n+t} \right) \prod_{i=0}^{v-1} y_i^{\sum_{j=1}^k s_{w_j-i}}$$

\square

Gale Robinson quivers have been specifically studied in depth already. A paper by Jeong, Musiker and Zhang [8] has a combinatorial interpretation for the coefficients in terms of brane tilings. As graphs admissible on surfaces, they also have combinatorial interpretations in terms of snake graphs and paths on marked surfaces. Furthermore a paper by Glick and Weyman [6] has an explicit formula for the coefficients formed by summing a function over order ideals of a poset. An advantage of our formula over the ones listed is that it is more explicit; it does not abstractly summing over a combinatorial object we do not know how to count, which is a pitfall of all the above formulas. Ours only involves understanding the partition function, which is simple when there are only two possible parts the number can be divided into.

5 Positivity of Exponents for Deformed F -Polynomials

In this section, we begin our study of deformed F -polynomials the deformed F -polynomial is, in fact, a polynomial.

Lemma 5.1. *Take a quiver Q and a sequence of mutations v_1, \dots, v_n . Every monomial in any F_n is expressible as the product of not necessarily distinct r_i with $i \leq n$.*

Proof. Assume there is a framed quiver Q and sequence of mutations v_1, v_2, \dots such that some F_n has a monomial not expressible as the product of the r_i 's. Take the smallest n for which this is true, and the let monomial of smallest degree in F_n for which this is true be m . Let V_{v_n} be the old label of vertex v_n . Let S_1 be the vertices corresponding to the set of base quiver vertices pointing in the same direction as the edges from frozen vertices. Let S_2 be the vertices corresponding to the other edges. Then, we have

$$F_n V_{v_n} = r_n \prod_{i \in S_1} V_i + \prod_{i \in S_2} V_i$$

It is easy to see that each F_i has a positive constant term of 1; namely V_{v_n} does, so then m is a term on the left hand side. Since there are no negative coefficients, the coefficient of m on the left hand side must be positive. However, every term on the right hand side is expressible as a product of r_i 's since every term in every F_i is, which is what the V_i 's are, and so is r_n . \square

Recall our set M_n for quiver Q and mutations v_1, \dots, v_n which is the set of fundamental r_i (r_i not expressible as the product of other r_j).

Lemma 5.2. *Every monomial in any F_n is expressible as the product of not necessarily distinct elements of M_n . In fact, it's expressible as the product of not necessarily distinct elements of M_m for any $m \geq n$, including M itself.*

Proof. By the previous lemma, we know every monomial is the product of r_i 's. Now, if an r_i is not fundamental, express it as a product of other r_i 's. Repeating this process must result in a product of fundamental r_i 's, since degrees are decreasing. \square

Lemma 5.3. *Given a framed quiver Q and a sequence of mutations v_1, \dots, v_n , $m \in M_n$, its coefficient can be computed as follows:*

Let g be the number of $j \leq n$ such that $m = r_j$ and r_j is green, and let r be the number of such $j \leq n$ so that r_j is red. Let $cf(m, V_i)$ be the coefficient of m in V_i . Then

$$-(g - r)C_n^{-1}(m) = \langle cf(m, V_1), cf(m, V_2), \dots, cf(m, V_v) \rangle$$

Proof. First, we compute the coefficient of m in F_t for any t . If m is fundamental, the only possible sequences of r_i 's producting to m consists of the individual r_i 's themselves that equal m . For all such r_i , plugging into our formula for F -polynomials gives that the inner expression $\phi(w_1, \dots, w_k)W(n, w_1, \dots, w_k)$ simplifies to $-C_{i,t}^{-1}[v_n, v_i]$. Since

$$-C_{i,t}^{-1}[v_n, v_i] = \delta(0, i)C_t^{-1}(m)[v_t],$$

summing over all the r_i equaling m gives $-(g-r)C_t^{-1}(m)[v_t]$. Now, we compute V_i . Take t so that t is the last step less than or equal to n where vertex v_t is mutated. If no occurrences of m as an r_i are between $t+1$ and n inclusive, then the coefficient of m is in fact $-(g-r)C_t^{-1}(m)[v_t]$, where the g and r correspond to number of occurrences of m preceding n . Note that

$$C_n^{-1} = A_n \cdots A_{t+1} C_t^{-1}$$

where each mutation A_i for $t+1 \leq i \leq n$ corresponds to a vertex v_i . It is easy to see that left multiplication by this matrix only affects the v_i th row of the matrix. As such, no mutation affects row v_t , so $C_t^{-1}(m)[v_t] = C_n^{-1}(m)[v_t]$.

Now we have the second case, where some occurrence of m proceeds spot t in the mutation sequence. Take a such occurrence of this to be at index s . Then, by the way the C matrix is defined, $C_s^{-1}(m)$ is the identity vector with a 1 at the spot $v_m \neq v_t$. Then, recalling that the v_t th row of the C^{-1} -matrix does not change between spot t and s and m , we have

$$C_t^{-1}(m)[v_t] = C_s^{-1}(m)[v_t] = C_n^{-1}(m)[v_t] = 0.$$

As such, the coefficient of m , which is $-(g-r)C_t^{-1}(m)[v_t]$ is 0, but this is also equal to $-(g-r)C_n^{-1}(m)[v_t]$ since this is also 0. Thus, in both cases, the expression simplifies to $-(g-r)C_n^{-1}(m)[v_t]$, and we are done. \square

Theorem 5.4. *Take a framed quiver Q and a mutation sequence v_1, \dots, v_n , with the following property: For any fundamental $m \in M_n$, the number of occurrences of $i \leq n$ such that $r_i = m$ and r_i is green is greater than the number where r_i is red. Then, S_n is a polynomial; that is none of the monomial terms have negative exponents if the r_n satisfy this property.*

Proof. Express any monomial in S_n as a product of fundamental terms. Choose a sequence of fundamentals m_1, \dots, m_k and express as $\prod m_i$. Since matrix operations are linear, we have that

$$-C_n^{-1} \left(\prod m_i \right) = \prod -C_n^{-1}(m_i).$$

We know that each monomial $-C_n^{-1}(m_i)$ has all positive exponents because of the following. Due to positivity of coefficients of F_n , every $m \in M_n$ is a valid monomial when acted on by $-C_n^{-1}$, since the coefficient vector multiplied by $(g-r)$ is the same as the exponent vector of $-C_n^{-1}(m_i)$, and the multiplication by $(g-r)$ preserves sign. \square

Corollary 5.5. *Given a framed quiver Q and a mutation sequence v_1, v_2, \dots with every r_i green, S_n is a polynomial.*

Proof. This is clear; for any r_i the number of green occurrences must be at least the number of red occurrences since there are only green occurrences. Thus, this follows from Theorem 5.4. \square

Almost conversely to Theorem 5.4, we have the following.

Corollary 5.6. *Take a framed quiver Q and a mutation sequence v_1, \dots, v_n , with the following property: there exists a fundamental m so that the number of occurrences of $i \leq n$ such that $r_i = m$ and r_i is red is greater than the number where r_i is green. Then, S_n is **not** a polynomial.*

Proof. Consider the monomial corresponding to m : $-C_n^{-1}(m)$. We claim it will have all negative exponents. Its coefficient vector in the original cluster of F -polynomials is $-(g-r)C_n^{-1}(m)$. This has to be always positive due to positivity of coefficients. Since $r > g$, this gives us that $-C_n^{-1}(m)$ has all negative entries, as desired. \square

Since we expect that the deformed F -polynomial is always a polynomial, we expect that the implied statement is always true.

Conjecture 5.7. *Given a framed quiver Q and a mutation sequence v_1, v_2, \dots , it is true that for any $m \in M$, for any n the number of occurrences of $i \leq n$ such that $r_i = m$ and r_i is green is at least the number where r_i is red.*

The reason we cannot get an if and only if statement with this method directly is that the case where $g = r$ for a fundamental monomial m is tricky to deal with. Replacing basic with fundamental provides an approach that circumvents this, though we do not know how to carry it through.

Conjecture 5.8. *Given a framed quiver Q and a sequence of mutations v_1, \dots, v_n , $p \in P_n$, the coefficient of p is the following:*

$$-C_n^{-1}(p) = \langle cf(p, V_1), cf(p, V_2), \dots, cf(p, V_v) \rangle$$

Theorem 5.9. *If Conjecture 5.8 is true, then for any framed quiver Q and mutation sequence v_1, \dots, v_n , S_n is a polynomial.*

Proof. For any basic monomial, the coefficient vector is the same as the exponent vector, by the conjecture. Any monomial can be rewritten as the product of basics. Each of these primes has positive coefficient vector, and therefore positive exponent vector, so the product will have positive exponent vector. \square

Conjecture 5.10. *Take a quiver Q and a sequence of mutations v_1, \dots, v_n . Let $\phi_{n,i}$ be the representation on the base quiver corresponding to the last step of the mutation process up to and including the n th step that occurred at vertex i . Let a monomial p be prime if any subrepresentation of $\phi_{n,i}$ with dimensions corresponding to the degree vector of p is indecomposable. Then the space of subrepresentations of $\phi_{n,i}$ with dimensions corresponding to the degree vector of p has Euler-Poincare characteristic of the quiver Grassmannian equal to $-C_n^{-1}(p)_i$*

Theorem 5.11. *If Conjecture 5.10 is true, then for any framed quiver Q and mutation sequence v_1, \dots, v_n , S_n is a polynomial.*

Proof. For any prime monomial, the coefficient vector is the same as the exponent vector, by the conjecture. Any monomial can be rewritten as the product of primes; rewrite it as the direct sum of two representations, and inductively decompose it into primes. Each of these primes has positive coefficient vector, and therefore positive exponent vector, so the product will have positive exponent vector. \square

6 Convergence in Periodic Case for Green Quivers

Lemma 6.1. *Take a framed quiver Q and a green mutation sequence v_1, \dots, v_n . For any r_i , we have that $-C_n^{-1}(r_i)$ is a monomial with positive exponents.*

Proof. We show each r_i is the product of fundamentals. Assume r_i is not a fundamental. Then, by definition, it is the product of other r_j 's. Repeating recursively, it is the product of fundamentals. Then, by a similar argument as before, $-C_n^{-1}(r_i)$ has to have all positive components, since $-C_n^{-1}$ of all its components in the decomposition into fundamentals is positive. \square

Lemma 6.2. *Given a green quiver Q and a mutation sequence v_1, v_2, \dots , all the r_i 's are distinct.*

Proof. Assume monomial m is both r_i and r_j for $j > i$. Consider the quiver after j mutation steps. First we show m is fundamental. If not, we will have

$$-C_j^{-1}(m) = \prod -C_j^{-1}(r_a).$$

However, the left side is just the 0 vector with a 1 at v_j , and this has degree 1 so it cannot be the product of other terms. Now we know that the coefficient vector of a fundamental r_i is simply $-(g-r)C_j^{-1}(r_i)$ which is 0 everywhere and 2 at v_i . In order for this to be true, at the previous step, the coefficient vector had to be 0 everywhere and -1 at v_i . This is because nothing other than v_i changed; the new r_i contributes 1, so the old term must contribute 1, and it was negated so must have been -1 before. This contradicts positivity of coefficients. \square

Lemma 6.3. *Take a green quiver Q , and a periodic mutation sequence $v_1, \dots, v_p, \dots, v_n$. For any monomial m in the stabilized polynomial, there exists a constant c so that for any sequence w_1, \dots, w_k $-C_n^{-1}(r_{w_1}) \cdots -C_n^{-1}(r_{w_k}) = m$, then $n - c \leq w_1, \dots, w_k$, independent of n .*

Proof. Consider instead the monomials after acted on by $-C_n^{-1}$. Let the sequence of mutations be $1, \dots, p, 1, \dots$, ending in a . We have

$$-C_n^{-1} = A_n \cdots A_1 = A_a \cdots A_1 A_p \cdots A_1 \cdots A_1.$$

For fixed a , consider each of the monomials $A_a \cdots A_1 \cdots (e_{v_a})$, at each possible termination point of the matrix product, and assume two of them are equal. Say the length of the sequences at these two termination points are b and c . Then take k larger than both b and c and equal to $a \pmod p$. Then

$$-C_{k-b,k}^{-1}(e_{v_a}) = -C_{k-c,k}^{-1}(e_{v_a})$$

However then

$$C(-C_{k-b,k}^{-1}(e_{v_a})) = C(-C_{k-c,k}^{-1}(e_{v_a})) \implies C_{k-b}(e_{v_{k-b}}) = C_{k-c}(e_{v_{k-c}})$$

This means both $r_{v_{k-b}}$ and $r_{v_{k-c}}$ are equal. However, we already showed that all the r_i are distinct in a green quiver. Thus, there is some k for any monomial p so that the only occurrence of p has to be at mutation $n - k$ after stabilizing the corresponding r_i , assuming $n \equiv a \pmod p$. Now, take the max of this value k (call it $\max(k)$) for all monomials $p|m$ over all the congruence classes $\pmod p$ (hence over all n). Since only monomials $p|m$ can contribute to m , we get that all contributors must be after $n - \max(k)$, as desired. \square

Theorem 6.4. *Given a green quiver Q , and a periodic mutation sequence v_1, \dots, v_p, \dots , for any i , S_i, S_{i+p}, \dots converges.*

Proof. Consider any monomial m and we show its coefficient converges. Take n larger than the c value for monomial m from the previous lemma. Thus, we have our equation:

$$S_n = \sum_{0 \leq w_1 \dots w_k \leq n-1} \phi(w_1, \dots, w_k) W(n, n - w_k, \dots, n - w_1) \prod_{i=0}^{v-1} y_i^{\sum_{j=1}^k \delta_{n-w_i, n} C_{n-w_i, n}^{-1}[j, v_n-w_i]}$$

which can be rewritten as

$$S_n = \sum_{0 \leq w_1 \dots w_k \leq c} \phi(w_1, \dots, w_k) W(n, n - w_k, \dots, n - w_1) \prod_{i=1}^v y_i^{\sum_{j=1}^k \delta_{n-w_j, n} C_{n-w_j, n}^{-1}[j, v_n-w_j]}$$

It would be clear these expressions were equal for sets of n where we could show $C_{n-w, n}^{-1}$ was only dependent w ; i.e. for a fixed w , the same regardless of n . However, this is a property true of all n in a congruence class $\pmod p$. Thus, the expression is in fact equal for all $n > c$. \square

Recall from before that stability was originally conjectured by Eager and Franco in their paper Color BPS, using the $dP1$ quiver. As a green quiver, our paper proves this case. We also write an explicit formula for the deformed F -polynomial in the proceeding section.

7 Exact Formulas for Deformed F -Polynomials

In this section, we present some exact formulas for these limit deformed F -polynomials we have now shown to exist. We restrict to the green quiver case, since that's where we have shown convergence.

7.1 Symmetric Quivers

For this subsection, we write the exact deformed F -polynomials for symmetric quivers. It does not illustrate much, but allows for simpler proofs in proceeding sections.

Theorem 7.1. *Let S_n be the deformed F -polynomial for the symmetric quiver Q . Then*

$$S_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_{v_n+1-i}^{\sum_{j=1}^k s_{n-w_j+1-i}}.$$

Proof. For this proof, consider vertices by their value $\pmod v$. Recall the formula from Theorem 4.3 which states

$$F_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_j+1-i}}.$$

We simply need to deform the exponents of the monomials. By Lemma 4.2,

$$r_k = \prod_{i=1}^v y_i^{s_{k+1-i}}$$

so the latter part of the expression can be rewritten as

$$\prod_{j=1}^k r_{w_j}.$$

Now we deform the exponents to get

$$\prod_{j=1}^k -C_n^{-1}(r_{w_j})$$

which is

$$\prod_{j=1}^k C_{w_j, n}^{-1}(e_{v_{w_j}}).$$

Now since the quiver is symmetric, each A_i is the same, with the vertices cyclic shifted according to v_i . Thus $C_{w_j, n}^{-1}$ will just be $A_{v_n} A_{v_n-1} \cdots A_{w_j}$ where there can be many periods in between. Mapping v_n to vertex 1, $v_n - 1$ to vertex 2 and so on, we get that $C_{w_j, n}^{-1}(e_{v_{w_j}})$ is $C_{0, n-w_j}(e_{v_{n+1-v_{w_j}}})$ under the new labeling. However, this is just r_{n-w_j} . Then we can use Lemma 4.2 to see that this is

$$r_{n-w_j} = \prod_{i=1}^v y_i^{s_{n-w_j+1-i}}.$$

Returning the vertices to their original labels, we get that each original r_k deforms to $\prod_{i=1}^v y_{v_n+1-i}^{s_{n-k+1-i}}$ and thus

$$S_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_{v_n+1-i}^{\sum_{j=1}^k s_{n-w_j+1-i}}.$$

□

7.2 $\widetilde{A}_{1,r}$

Definition 7.2. The quiver $\widetilde{A}_{1,r}$ is on vertices $1, \dots, r+1$ with the following edges:

- An edge from $i \rightarrow i+1$ for $1 \leq i \leq r$.
- An edge from $r+1 \rightarrow 1$.

In this section, we look for explicit formulas for deformed F -polynomials for the quiver $\widetilde{A}_{1,r}$. In her paper [18], Grace Zhang looks at the quiver $\widetilde{A}_{1,1}$. She shows the following. Let S_∞ be the deformed F -polynomial for the quiver $\widetilde{A}_{1,2}$. Then, up to swapping y_1 and y_2 , we have

$$S_\infty = 1 + \frac{y_1}{(1 - y_1 y_2)^2}.$$

Our theorem generalizes this result.

Theorem 7.3. Let S_∞ be the deformed F -polynomial for the quiver $\widetilde{A}_{1,r}$. Then

$$S_\infty = 1 + \frac{y_{r+1}(1 + y_r + y_r y_{r-1} + \dots + y_r \cdots y_2)}{(1 - y_1 \cdots y_{r+1})^2}$$

up to cyclic shift of variables.

Proof. Let W be the set of sequences $1 \leq w_1 \leq \dots \leq w_k \leq n$. By Theorem 7.1 we know that

$$S_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_{v_n+1-i}^{\sum_{j=1}^k s_{n-w_j+1-i}}$$

where $s_i = \lfloor \frac{i}{r} \rfloor$.

Consider any monomial m that we are trying to find the coefficient of in the stabilized polynomial which has nonzero coefficient. We claim that the exponent of y_{v_n} is one greater than the exponent of y_{v_n+1} in any term m with nonzero coefficient. Because \mathbf{s} increases every r terms and there are $r+1$ vertices, the difference between the exponent of y_{v_n} and y_{v_n+1} is going to be k . This also means that all sequences creating a given monomial are all the same length. Thus, we just have to show that the only terms with nonzero coefficient in the limit are the ones with all corresponding sequences \mathbf{w} having length 1. Assume some sequence \mathbf{w} with length greater than 1 corresponds to some monomial which has nonzero coefficient in the limit, specifically for S_n and all future $S_{n+k(r+1)}$. In S_n it corresponds to w_1, \dots, w_k . Then consider the polynomial $S_{n+n(r+1)}$. The sequences $w_1 + n(r+1), \dots, w_k + n(r+1)$ would contribute to this monomial in $S_{n+n(r+1)}$. However, in the original un-deformed polynomial, this would mean the corresponding sequence $r_{w_1+n(r+1)} \cdots r_{w_k+n(r+1)}$ products to the corresponding term. However, this terms clearly has degree too large in the original polynomial, so must have 0 coefficient. Therefore, it suffices to sum over all sequences \mathbf{w} with length 1. Then we get

$$S_\infty = \sum_{1 \leq w \leq n} s_{n-w} \prod_{i=1}^v y_{v_n+1-i}^{s_{n-w+1-i}}$$

where $s_i = \lfloor \frac{i}{r} \rfloor$. This simplifies to the following. Let W be the set of sequences $\lfloor \frac{n}{r} \rfloor \geq w_{v_n} \geq \dots \geq w_{v_n+1} \geq 0$ where w_{v_n} is one more than w_{v_n+1} . Then taking limit over sequences that end in the same mutation step v_n ,

$$S_\infty = \sum_{\mathbf{w} \in W} w_{v_n} \prod_{i=1}^{r+1} y_{v_n+1-i}^{w_{v_n}}$$

This can be seen to rewrite to

$$S_\infty = 1 + \frac{y_{r+1}(1 + y_r + y_r y_{r-1} + \dots + y_r \cdots y_2)}{(1 - y_1 \cdots y_{r+1})^2}$$

up to cyclic shift of variables, dependent on the last mutation step. \square

7.3 K_r

In this section, we investigate the deformed F -polynomial for the K_r quiver. It is worthwhile to note that $K_2 = \widehat{A}_{1,1}$. As such, this case can be viewed as a different generalization of the Zhang's result for the $\widehat{A}_{1,1}$ quiver.

Definition 7.4. Define the sequence s as $\dots, s_{-2} = s_{-1} = 0, s_0 = 1$ and $s_{k+2} = ns_{k+1} - s_k$ for $k \geq 0$. Let $p = \frac{r+\sqrt{r^2-4}}{2}$ Let the norm at b of a sequence $1 \leq w_1 \leq \dots \leq w_k$ be

$$N_b(\mathbf{w}) = \sum_{i=1}^k b^{w_i}.$$

The explicit value of any given s_i for $i \geq 0$ can be calculated as follows:

$$s_i = \frac{1}{\sqrt{r^2-4}} p^i - \frac{1}{\sqrt{r^2-4}} p^{-i}.$$

Theorem 7.5. *The deformed F -polynomial S_∞ is calculated as follows. Let Q be the set of sequences \mathbf{w} with $0 \leq w_1 \leq \dots \leq w_k$ with $N_{1/p}(\mathbf{w})$ at most 1.*

$$S_\infty = 1 + \sum_{\mathbf{w} \in W} \phi(\mathbf{w}) \prod_{i=1}^k \left(s_{w_i} - \sum_{j=1}^i s_{w_i-w_j} + s_{w_i-w_{j-2}} \right) y_1^{\sum_{i=1}^k s_{w_i}} y_2^{\sum_{i=1}^k s_{w_i-1}}$$

Alternately,

$$S_\infty = 1 + \sum_{\mathbf{w} \in W} \phi(\mathbf{w}) \prod_{i=1}^k \left(s_{w_i} - \sum_{j=1}^i s_{w_i-w_j} + s_{w_i-w_{j-2}} \right) y_1^{\left\lfloor \frac{1}{\sqrt{r^2-4}} p N_p(\mathbf{w}) \right\rfloor} y_2^{\left\lfloor \frac{1}{\sqrt{r^2-4}} N_p(\mathbf{w}) \right\rfloor}$$

up to a permutation of variables.

Proof. Recall the statement of our formula from Theorem 7.1:

$$S_n = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \left(\prod_{i=1}^k s_{n-w_i} + \sum_{j=i}^k -s_{w_j-w_i} + s'_{w_j-w_i} \right) \prod_{i=1}^v y_{v_{n+1-i}}^{\sum_{j=1}^k s_{n-w_j+1-i}}.$$

Noting that there are only two vertices, this simplifies to

$$S_n = \sum_{1 \leq w_1 \dots w_k \leq n} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{w_i}) - \sum_{j=1}^{i-1} (s_{w_j-w_i} + s_{w_j-w_i-2}) \right) y_1^{\sum_{j=1}^k s_{w_i}} y_2^{\sum_{j=1}^k s_{w_i-1}}$$

where y_1 and y_2 are swapped if n is even. We ignore this distinction in the proof that follows. This is exactly our desired expression, except we need to limit the set of sequences we are summing over, using the norm condition. We take a monomial m with degree in y_1 being d . Then, all possible sequences w that allow for this monomial have at most d terms and all $w_i < d$ (d is a very crude bound, but suffices). Take n extremely large. Now, we look at the degree of the term corresponding to w in the original F -polynomial. We want to show both that if the norm is at most 1, the degree will be this must be less than or equal to s_{n-1} , and if norm is greater than 1 it will be larger. The degree is

$$\sum_{i=1}^k s_{n-w_i} = \frac{1}{\sqrt{r^2-4}} (p^n N_{1/p}(w) - \sum_{i=1}^k (1/p)^{n-w_i})$$

by using the explicit form of the s_i . s_{n-1} can be expressed as

$$\frac{1}{\sqrt{r^2-4}}p^n - \frac{1}{\sqrt{r^2-4}}p^n$$

When n goes to infinity, the first term of each expression dominates, and this easily proves the desired result; the first is smaller when norm is less than 1, and larger otherwise. If the norm is exactly 1, the first terms of each expression are equal, so we compare the second term. The first expression clearly has a smaller first term, so the result holds in this case as well.

Now we prove the second part; that

$$\sum_{j=1}^k s_{w_j} = \left\lfloor \frac{1}{\sqrt{r^2-4}} N_p(q) \right\rfloor$$

and

$$\sum_{j=1}^k s_{w_{j-1}} = \left\lfloor \frac{1}{\sqrt{r^2-4}} p N_p(q) \right\rfloor$$

for a sequence with norm at most 1. For the first part, first rewrite

$$\sum_{j=1}^k s_{w_j} = \frac{1}{\sqrt{r^2-4}} (p N_p(w) - p^{-1} N_{1/p}(w))$$

The second term (after expanding) is strictly greater than 0 and less than 1, and since the expression must be an integer, we have proven our desired result. We have

$$\sum_{j=1}^k s_{w_j} = \frac{1}{p\sqrt{r^2-4}} (N_p(w) - N_{1/p}(w))$$

Again the second term after expanding has to be greater than 0 and less than 1, proving the desired result. \square

Corollary 7.6. *For the deformed F -polynomial S_∞ of the quiver K_r , the terms that may have nonzero coefficient are exactly the ones of the form $y_1^{\lfloor pa \rfloor + 1} y_1^a$ for integers $a \geq 0$. This coefficient is calculated as follows. Let W be the set of sequences \mathbf{w} with $0 \leq w_1 \leq \dots \leq w_k$ and $\sum_{\mathbf{w} \in W} s_{w_i-1} = a$*

$$\phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{w_i}) - \sum_{j=1}^{i-1} (s_{w_j-w_i} + s_{w_j-w_i-2}) \right)$$

Proof. We need to find

$$\left\lfloor p \left(\left\lfloor \frac{1}{\sqrt{r^2-4}} N_p(q) \right\rfloor \right) \right\rfloor + 1$$

Recall from the previous proof that our expression of $\left\lfloor \frac{1}{\sqrt{r^2-4}} N_p(q) \right\rfloor$ is also

$$\frac{1}{\sqrt{r^2-4}} N_p(q) - \frac{1}{\sqrt{r^2-4}} N_{1/p}(q)$$

Multiplying by p gives

$$\frac{p}{\sqrt{r^2-4}} N_p(q) + \frac{p}{\sqrt{r^2-4}} N_{1/p}(q)$$

which is

$$\sum_{i=1}^{\lfloor q \rfloor} s_{w_i} + \frac{1}{p\sqrt{r^2-4}} N_{1/p}(\mathbf{w}) - \frac{1}{\sqrt{r^2-4}} N_{1/p}(\mathbf{w})$$

which is

$$\sum_{i=1}^{|q|} s_{w_i} + \frac{p-1}{p} \frac{1}{\sqrt{r^2-4}} N_{1/p}(q)$$

The second term is non-strictly less than 1, but greater than 0, so taking the floor and adding 1 gives $\text{sum}_{i=1}^{|q|} s_{w_i}$ as desired for the first part.

For the second part, we know that we get a given term by summing the expression

$$\phi(w_1, \dots, w_k) \prod_{i=1}^k \left((s_{w_i}) - \sum_{j=1}^{i-1} (s_{w_j-w_i} + s_{w_j-w_i-2}) \right)$$

over sequences \mathbf{w} with $0 \leq w_1 \leq \dots \leq w_k$ with $\sum_{\mathbf{w} \in W} s_{w_{i-1}} = a$. Thus, it suffices to prove that $a = \left\lfloor \frac{p}{\sqrt{r^2-4}} N_p(q) \right\rfloor$. Note that

$$s_{w_{i-1}} = \frac{1}{p\sqrt{r^2-4}} N_p(\mathbf{w}) - \frac{p}{\sqrt{r^2-4}} N_{1/p}(\mathbf{w}).$$

The second term is less than 1, so we get that the overall expression is just the floor of the first part, as desired. \square

The deformed F -polynomial is interesting for two reasons: it allows for the convenient norm property to determine which monomials are included. Further, the renormalization has y_1 and y_2 uniquely determine each other.

7.4 Gale Robinson Quivers

We now look at the explicit deformed F -polynomial for the Gale Robinson Quivers. Essentially we just plug into our explicit formula and nothing extremely interesting happens, but it allows us to completely close the Eager and Franco's original conjecture data around the $dP1$ quiver.

Theorem 7.7. *Take the quiver $G_{v,r,t}$. Define a sequence \mathbf{s} with s_i as the number of partitions of i into parts $r, v-r$ (0 for negative values of i , 1 for 0). Let W be the set of sequences \mathbf{w} with $0 \leq w_1 \leq \dots \leq w_k$.*

$$S_\infty = \sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left(s_{w_i} + \sum_{j=1}^{i-1} (-s_{w_j-w_i} - s_{w_j-w_i-v} + s_{w_j-w_i-t} + s_{q_j-q_i-n+t}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_i-j+1}}$$

Proof. We have by Theorem 7.1 that

$$S_n = \sum_{0 \leq w_1 \dots w_k \leq n-1} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left(s_{w_i} + \sum_{j=1}^{i-1} (-s_{w_j-w_i} - s_{w_j-w_i-v} + s_{w_j-w_i-t} + s_{w_j-q_i-n+t}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_i-j+1}}$$

since that is how to calculate the corresponding sequences, and so taking the limit gives

$$S_\infty = \sum_{0 \leq w_1 \dots w_k} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left(s_{w_i} + \sum_{j=1}^{i-1} (-s_{w_j-w_i} - s_{w_j-w_i-v} + s_{w_j-w_i-t} + s_{w_j-q_i-n+t}) \right) \prod_{i=1}^v y_i^{\sum_{j=1}^k s_{w_i-j+1}}$$

\square

Now we find the exact deformed F -polynomial for the quiver $dP1 = G_{4,2,1}$. If we can, it is always more interesting to calculate the coefficient of a given monomial directly, rather than summing up a number of terms that could possibly be contributing to the same polynomial. We will do this for the $dP1$ quiver, since the sequence \mathbf{s} is easy to understand.

Corollary 7.8. *Take the quiver dP1 and mutation sequence $1, 2, 3, 4, 1, 2, \dots$ which has n terms. The coefficient of the term $y_1^a y_2^b y_3^c y_4^d$ is calculated as follows. Let W be the set of sequences of nonnegative integers and half-integers so that*

- $a - c$ terms are integers.
- $b - d$ terms are half-integers.
- The sum of all the integer terms is c .
- The sum of the floors of all the half integer terms is d .

Then the desired coefficient is

$$\sum_{\mathbf{w} \in W} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left(\frac{(-1)^{2w_i} + 1}{2} (w_i + 1) - \sum_{j=1}^{i-1} (-1)^{2w_j - 2w_i} (2w_j - 2w_i + 4) \right)$$

Proof. Let $s(k)$ be k if k is a positive integer, and 0 otherwise. Noting that s_k as defined in the Gale Robinson quiver is $\frac{k}{2}$ if k is even and 0 otherwise, plugging in gives:

$$S_\infty = \sum_{0 \leq w_1 \dots w_k, w_i} \phi(w_1, \dots, w_k) \prod_{i=1}^k \left(\frac{((-1)^{w_i} + 1)w_i}{4} - \sum_{j=1}^{i-1} (-1)^{w_j - w_i} (w_j - w_i + 4) \right) \prod_{i=1}^4 y_i^{\sum_{j=1}^k \frac{(-1)^{w_i - j + 1} + 1}{4}}$$

Letting $q_i = \frac{w_i}{2}$ gives the following expression.

$$S_\infty = \sum_{q \in Q} \phi(q_1, \dots, q_k) \prod_{i=1}^k \left(q(w_i) - \sum_{j=1}^{i-1} (-1)^{2q_j - 2q_i} (2q_j - 2q_i + 4) \right) \prod_{i=1}^4 y_i^{\sum_{j=1}^k s(q_j - 0.5i)}$$

For any term $y_1^a y_2^b y_3^c y_4^d$ it is contributed to by the sequences described in the corollary statement. Adding the contributions proves the desired result. \square

8 Future Research

8.1 The Formula

Future research involving our formula in Theorem 3.1 mostly consists of finding more use cases for the formula. For example, it might be possible to group terms and use the formula to recover positivity of coefficients, at least in certain cases. We might be able to do this by grouping terms of the formula.

Secondly, we know that the Laurent Phenomenon proves that the F -polynomials are in fact polynomials. However, our formula has nontrivial expansion for the coefficient of all monomials. Thus, for monomials that are too large, or obviously otherwise have coefficient 0, we get an expression involving the C -matrix entries that must be zero. We have not yet investigated these; it might be interesting. It might even help with positivity; one might be able to say that if large terms have coefficient 0, the smaller terms must have coefficient positive. The following provides intuition for why such a proof might work.

Lemma 8.1. *Take a quiver Q and a mutation sequence v_1, \dots, v_n . If each r_i^k can only be constructed as a product of r_i 's, and not any of the other terms, then the coefficient of r_i must be positive.*

Proof. The coefficient of r_i by our formula is $a_{i,n}$, and the coefficient of r_i^k is

$$\frac{1}{k!} \prod_{i=1}^k (a_{i,n} - \sum_{j=1}^{k-1} (a_{i,j} - b_{i,j}))$$

which is

$$\frac{1}{k!} \prod_{i=1}^k (a_{i,n} - k + 1)$$

This is only 0 for large k when $a_{i,n}$ is positive. □

This idea is complicated to generalize, since in large enough monomials, every part-contribution (the particular sequence we are summing over) need not be 0. However, there might be a clever way to group terms to deal with this.

Another possible avenue of research would be to see how our work relates to dilogarithm identities. Keller [9] and Nakanashi [16] discuss these identities, and they appear to be an area where explicit formulas may allow for better results.

8.2 Deformed F -Polynomials

Regarding deformed F -polynomials, we have the following three conjectures.

Conjecture 8.2. *Given a framed quiver Q and a mutation sequence v_1, v_2, \dots, S_n is actually a polynomial; that is none of the monomial terms have negative exponents.*

Conjecture 8.3. *Given a framed quiver Q and a mutation sequence v_1, v_2, \dots, S_n is bounded; i.e. for each monomial term $\prod x_i^{a_i}$, there is some $c(a_1, \dots, a_n)$ dependent on Q and the mutation sequence so that the coefficient of $\prod x_i^{a_i}$ in S_n is less than $c(a_1, \dots, a_n)$.*

Conjecture 8.4. *Given a framed quiver Q and a periodic mutation sequence $v_1, \dots, v_p, v_1, \dots, v_p, \dots$, such that performing the mutations v_1, \dots, v_p on the base quiver returns the quiver back to the original, for every i , the sequence $S_i, S_{i+p}, S_{i+2p}, \dots$ converges.*

We proved the first and third in the case of green quivers. The first is very related to the coefficients in our formula being positive. Recall that our proof of positivity in the green case relied on positivity of coefficients. It might be possible to generalize this, using our formula to understand the coefficient of “basic” terms via the formula. This would in turn prove Conjecture 5.8, thus proving the result. We might also be able to achieve a direct proof using a concept like in lemma 8.1.

We have not spent much time investigating the second conjecture. We need to extend the proof used for convergence in the periodic green case in two ways. We need to remove the condition periodic and the condition green. It would be especially interesting and probably doable to remove the periodic condition.

For the third conjecture, again, we have proven it in the case of green quivers. The arguments do not extend directly from the green case, though a proof is likely similar to a full proof of the previous conjectures.

9 Acknowledgments

This research was carried out as part of the 2018 Mathematics REU at University of Minnesota, Twin Cities. It was supported by the NSF RTG grant DMS-1745638. The author would like to thank mentor Gregg Musiker for his invaluable guidance and support, as well as TA Esther Banaian.

References

- [1] I. Assem, G. Dupont, R. Schiffler, and D. Smith. Friezes, Strings and Cluster Variables. *ArXiv e-prints*, September 2010.
- [2] H. Derksen, J. Weyman, and A. Zelevinsky. Quivers with potentials and their representations I: Mutations. *ArXiv e-prints*, April 2007.

- [3] R. Eager and S. Franco. Colored BPS pyramid partition functions, quivers and cluster transformations. *Journal of High Energy Physics*, 9:38, September 2012.
- [4] S. Fomin and A. Zelevinsky. Cluster algebras I: Foundations. *ArXiv Mathematics e-prints*, April 2001.
- [5] S. Fomin and A. Zelevinsky. Cluster algebras IV: Coefficients. *ArXiv Mathematics e-prints*, February 2006.
- [6] M. Glick and J. Weyman. Gale-Robinson quivers: from representations to combinatorial formulas. *ArXiv e-prints*, October 2017.
- [7] M. Gross, P. Hacking, S. Keel, and M. Kontsevich. Canonical bases for cluster algebras. *ArXiv e-prints*, November 2014.
- [8] In-Jee Jeong, Gregg Musiker, and Sicong Zhang. Gale-robinson sequences and brane tilings. 01 2013.
- [9] B. Keller. On cluster theory and quantum dilogarithm identities. *ArXiv e-prints*, February 2011.
- [10] B. Keller. Cluster algebras and derived categories. *ArXiv e-prints*, February 2012.
- [11] K. Lee and R. Schiffler. Positivity for cluster algebras. *ArXiv e-prints*, June 2013.
- [12] Kyungyong Lee. On cluster variables of rank two acyclic cluster algebras. *Annals of Combinatorics*, 16(2):305–317, Jun 2012.
- [13] Kyungyong Lee and Ralf Schiffler. A combinatorial formula for rank 2 cluster variables. *Journal of Algebraic Combinatorics*, 37(1):67–85, Feb 2013.
- [14] G. Musiker, R. Schiffler, and L. Williams. Positivity for cluster algebras from surfaces. *ArXiv e-prints*, June 2009.
- [15] Gregg Musiker and Ralf Schiffler. Cluster expansion formulas and perfect matchings. *Journal of Algebraic Combinatorics*, 32(2):187–209, Sep 2010.
- [16] T. Nakanishi. Tropicalization method in cluster algebras. *ArXiv e-prints*, October 2011.
- [17] R. Schiffler and H. Thomas. On cluster algebras arising from unpunctured surfaces. *ArXiv e-prints*, December 2007.
- [18] G. Zhang. Stable Cluster Variables. *University of Minnesota, Twin Cities REU Report*, 2016.