

Theory of Ordinary Differential Equations

Arnd Scheel, VinH 509, scheel@umn.edu

— *Optional exercises* —

(i) An exercise in abstraction:

(a) Show that $C^0([0, 1])$ is a Banach space when equipped with the norm

$$\|u\|_{C^0} = \sup_{t \in [0, 1]} |u(t)|.$$

(b) Show that $C^1([0, 1]) \cap \{u | u(0) = 0\}$ is a Banach space when equipped with the norm

$$\|u\|_{C^1} = \sup_{t \in [0, 1]} (|u'(t)| + |u(t)|).$$

(c) Show that the operator

$$L : X \rightarrow Y, \quad u \mapsto u',$$

is linear and bounded.

(d) Show that L is bounded invertible and describe the inverse.

(ii) Find the solution to $x' = Ax \in \mathbb{R}^2$, $A = \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}$, with $x(0) = (0, 1)^T$, for any $\mu \in \mathbb{R}$.

(ii) Find the solution to $x' = Ax \in \mathbb{R}^2$, $A = \begin{pmatrix} 1 & \mu \\ 0 & 1 \end{pmatrix}$, with $x(0) = (0, 1)^T$, for any $\mu \in \mathbb{R}$.