

Homework 4

Let E, F, G denote Banach spaces.

1. Let $T \in \mathcal{L}(E)$ be a linear continuous operator on a complex Banach space E . By $\mathcal{R} : \rho(T) \rightarrow \mathcal{L}(E)$ we denote the resolvent function of T . Prove that:

- (i) $\mathcal{R}(\lambda_1) - \mathcal{R}(\lambda_2) = (\lambda_1 - \lambda_2)\mathcal{R}(\lambda_1)\mathcal{R}(\lambda_2), \quad \forall \lambda_1, \lambda_2 \in \rho(T)$
- (ii) \mathcal{R} is differentiable on $\rho(T)$ (find its derivative)
- (iii) $\lim_{|\lambda| \rightarrow \infty} \mathcal{R}(\lambda) = 0$
- (iv) the spectrum $\sigma(T)$ is always nonempty
- (v) if $\lambda \in \sigma(T)$ then $\lambda^n \in \sigma(T^n)$
- (vi) the spectral radius: $\max\{|\lambda|; \lambda \in \sigma(T)\} = \lim_{n \rightarrow \infty} \|T^n\|^{1/n}$.

2. Let $T \in \mathcal{K}(E)$. Is $\text{Range}(T)$ always separable?

3. Prove that $T \in \mathcal{L}(E, F)$ is compact iff

$$T^* : (\bar{B}_{F^*}, \text{weak}^*) \rightarrow E^*$$

is continuous.

4. (i) Let $T \in \mathcal{K}(E, F)$ and let $S \in \mathcal{L}(F, G)$ be injective. Prove that for every $\epsilon > 0$ there exists $C > 0$ so that:

$$\|Tx\| \leq \epsilon\|x\| + C\|STx\| \quad \forall x \in E.$$

(ii) Let Ω be a ball in \mathbf{R}^n . Prove that for every $\epsilon > 0$ there exists $C > 0$ so that:

$$\|\nabla u\| \leq \epsilon\|D^2u\| + C\|u\| \quad \forall u \in \mathcal{C}^2(\bar{\Omega}),$$

where $\|\cdot\|$ denotes the $\mathcal{C}^0(\bar{\Omega})$ norm of a given function.

(iii) Prove that in (ii) one can take $C = 4(\min(\epsilon, 1/R))^{-1}$, where R is the radius of the ball Ω .

5. Give an example of a compact operator on l_2 , whose point spectrum is empty.

6. Let T be the integral Hilbert-Schmidt operator, as in Problem 5 Homework 2. Prove that:

$$\dim \text{Ker}(\text{Id} - T) \leq \|K\|_{L^2(\Omega \times \Omega)}^2.$$