

Date due: October 16, 2006. Either hand it to me in class or put it in my mailbox by 3:30.

1. Use GAP to show that

$$\langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^2 = (bc)^3 = (ca)^5 = 1 \rangle \cong A_5 \times C_2.$$

2. The generalized quaternion group of order  $2^n$  has a presentation

$$\langle a, b \mid a^{2^{n-1}} = 1, b^2 = a^{2^{n-2}}, bab^{-1} = a^{-1} \rangle.$$

Use GAP to investigate the generalized quaternion group of order 32. Get a list of the orders of the elements. Compute the derived subgroup and the center. Draw a picture of the lattice of subgroups of this group. What is the minimum degree of a faithful permutation representation of this group?

3. (a) Show that every homomorphism of  $G$ -sets  $\Omega \rightarrow \Psi$  where  $\Psi$  is transitive is necessarily an epimorphism.  
 (b) Let  $\Psi$  be a transitive  $G$ -set. Show that every  $G$ -set mapping  $\Psi \rightarrow \Psi$  is a bijection. (It is not assumed that  $G$  be finite here.)  
 (c) Let  $H$  and  $K$  be subgroups of  $G$ . Show that every homomorphism of  $G$ -sets  $G/H \rightarrow G/K$  is a composite  $G/H \rightarrow G/J \rightarrow G/K$  where  $H \leq J$ ,  $J$  is conjugate to  $K$ , and the mapping  $G/H \rightarrow G/J$  is  $xH \mapsto xJ$ .
4. Use GAP to show that the group  $\langle (1, 5)(2, 6), (1, 3)(4, 6), (2, 3)(4, 5) \rangle$  is isomorphic to  $S_4$ .
5. (Question 12.5 from the handout) A group  $G$  is *injective*  $\Leftrightarrow$  whenever we are given a subgroup  $A$  of a group  $B$  and a homomorphism  $f : A \rightarrow G$  there exists a homomorphism  $g : B \rightarrow G$  so that the restriction of  $g$  to  $A$  is  $f$ . Prove that injective groups have order 1. [Hint (D.L. Johnson): let  $A$  be free on  $\{a, b\}$  and let  $B = A \rtimes \langle c \rangle$  where  $c$  has order 2 and  $cac^{-1} = b, cbc^{-1} = a$ .]
6. (Question 12.7 from the handout) Let  $X = \{x_k \mid k \in K\}$  and let  $Y \subseteq X$ . If  $F$  is free on  $X$  and  $H$  is the normal subgroup generated by  $Y$ , show that  $F/H$  is free.
7. (Question 12.8 from the handout) Show that a free group  $F$  on  $\{x, y\}$  has an automorphism  $f$  with  $f(f(a)) = a$  for all  $a \in F$  and with the further property that  $f(a) = a$  if and only if  $a = 1$ .

**Extra Questions: do not hand in**

8. Use GAP to show that  $SL(2, 5)$  has a normal subgroup of order 2 such that the quotient is isomorphic to  $A_5$ . Show that  $SL(2, 5)$  has no subgroup isomorphic to  $A_5$ . Identify the Sylow 2-subgroups of  $SL(2, 5)$ .

9. Use GAP to investigate the groups

$$\langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^2 = (bc)^3 = (ac)^4 = 1 \rangle$$

and

$$\langle a, b, c \mid a^2 = b^2 = c^2 = (ab)^2 = (bc)^3 = (ac)^3 = 1 \rangle$$

In each case identify the quotient by the center  $G/Z(G)$  and determine whether or not  $G = Z(G) \times H$  for some subgroup  $H$ .